



中国科学院科学出版基金资助出版

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

月刊 SCI 收录 2010年7月 第31卷 第7期



目次

综述

717 (特约)

新型微纤结构催化/吸附填料研究进展
凌敏, 赵国锋, 曹发海, 路勇

研究快讯

725 (国际版)

AlCl₃/Al-MCM-41 催化剂上萘与丙烯的烷基化反应
唐华, 纪敏, 王新葵, 贺民, 蔡天锡

研究论文

729 (国际版/英文)

预吸附吡啶增强二甲醚在丝光沸石上羰基化反应的稳定性
刘俊龙, 薛会福, 黄秀敏, 吴培豪, 黄信灵, 刘尚斌, 申文杰

739 (国际版/英文)

钼硫碘纳米线结构特性和化学改性的理论研究
温书豪, 侯柱锋, 刘建勇, 何国钟

747 (国际版/英文)

LaZn_xFe_{1-x}O₃ 钙钛矿型纳米催化剂的制备、表征及其甲苯燃烧性能
Seyed Ali HOSSEINI, Mohammad Taghi SADEGHI, Abdolali ALEMI, Aligholi NIAEI, Dariush SALARI, Leila KAFI-AHMADI

751 (国际版/英文/封面文章)

紫外光照射下 Fe-碳纳米管/TiO₂ 复合材料降解罗丹明 B
ZHANG Kan, MENG Zeda, OH Wonchun

759 (国际版/英文)

H₃PO₄/MCM-41 催化苯酚与叔丁醇气相烷基化反应
Mehran GHIACI, Behzad AGHABARARI

765 (国际版)

水合碱金属卤化物催化 CO₂ 与环氧化物合成环状碳酸酯
周喜, 张毅, 杨先贵, 姚洁, 王公应

769 (国际版/英文)

温和条件下 Cu-Zn/Al₂O₃ 催化剂上酯加氢制醇
袁鹏, 刘仲毅, 张万庆, 孙海杰, 刘寿长

776

离子液中磷酸铝方英石的低温合成及表征
徐仁顺, 张维萍, 韩秀文, 包信和

781

Rh-Au/γ-Al₂O₃ 三效纳米催化剂的制备与表征
刘立成, 訾学红, 戴洪兴, 赵震, 王新平, 何洪

788

ZSM-35 分子筛的静态合成和晶化机理
鞠晓花, 范峰滔, 田福平, 冯兆池

793

ZrO₂ 对 CuMn₂/Al-Ti 整体式催化剂催化苯燃烧反应性能的影响
张志强, 贺站锋, 王娟芸, 王奕祎, 李吉花, 蒋毅, 陈君和, 张小霞

797

Fe-Ni 共掺杂 ZnO 的制备及其光催化降解甲基橙活性
傅天华, 高倩倩, 刘斐, 代华均, 寇兴明

803

以四乙基氢氧化铵-四甲基氯化铵为模板剂水热法合成 UZM-5 沸石
刘艳, 李英霞, 李建伟, 陈标华

809

Cr 助剂对 Fe 基催化剂结构和费托合成性能的影响
王虎林, 杨勇, 王洪, 青明, 相宏伟, 李永旺

817

介质阻挡放电与 CuZSM-5 结合方式对脱除 NO_x 的影响
陈刚, 孙琪, 石雷, 牛金海, 宋志民

822

固体磺酸促进亚硝酸异戊酯催化氧化苯甲醇反应
盛学斌, 马红, 李德财, 何静, 徐杰

827

SBA-15 固载离子液体功能化脯氨酸的制备及其催化 Knoevenagel 缩合反应
沈加春, 郭建平, 孙艳美, 唐斌艳, 陈小花, 尹笃林

833

以催化裂化废催化剂粉合成超细 Y 型分子筛
刘欣梅, 梁海宁, 李亮, 杨婷婷, 阎子峰

839

Au/SBA-15 的制备及其催化 CO 氧化反应性能
苏继新, 张慎平, 马丽媛, 屈文, 张明博

846

SiO₂ 负载壳聚糖席夫碱钨催化剂催化芳香醛酮还原为芳香烃
贺海峰, 龚树文, 刘丽君, 崔庆新, 尹汉东

851

无氯 Cu/AC 催化剂的制备及其催化气相甲醇氧化羰基化反应性能
王瑞玉, 李忠, 郑华艳, 谢克昌

相关信息

838 关于召开现代催化研究方法讲习班的通知

857 作者索引



Supported by the Science Publication
Foundation of the CAS

催化学报
(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

Monthly Vol. 31 No. 7 July 2010



Contents

Review


Chin. J. Catal., 2010, 31: 717–724 doi: 10.3724/SP.J.1088.2010.00146

Advances in Research on Novel Microfibrous Structured Catalytic/Adsorbent Packings

LING Min, ZHAO Guofeng, CAO Fahai, LU Yong*

East China Normal University; East China University of Science and Technology

Microfibrous structure



Beneficial properties

- Unique form factors
- High contacting efficiency
- Enhanced mass/heat transfer
- Low pressure drop compared to randomly packed beds
- Uniform residence time
- Avoiding separation issues (e.g., suspension reactors)
- Facilitation of scale-up
- Ease of functionalization

Research practices

- Miniature H₂ generation and cleanup
- High efficiency air filtration
- Electrochemical energy storage
- Selective oxidation of alcohols
- F-T synthesis
- ...

Process Intensification

New-type microfibrous structured packings demonstrate a beneficial manner different from traditional contacting schemes such as packed beds. The latest progress in their use toward process intensification has been reviewed.


Communication

Chin. J. Catal., 2010, 31: 725–728 doi: 10.1016/S1872-2067(09)60080-2

Alkylation of Naphthalene with Propylene Catalyzed by Aluminum Chloride Immobilized on Al-MCM-41

TANG Hua, JI Min*, WANG Xinkui, HE Min, CAI Tianxi

Dalian University of Technology




Alkylation of naphthalene with propylene can be carried out over $\text{AlCl}_3/\text{Al-MCM-41}$ catalysts in a slurry bubble column reactor at $85\text{ }^\circ\text{C}$. The pore diameter influences the conversion of naphthalene and selectivity for 2,6-diisopropylnaphthalene.

Chin. J. Catal., 2010, 31: 729–738 doi: 10.1016/S1872-2067(09)60081-4

Stability Enhancement of H-Mordenite in Dimethyl Ether Carbonylation to Methyl Acetate by Pre-adsorption of Pyridine

LIU Junlong, XUE Huifu, HUANG Xiumin, WU Pei-Hao, HUANG Shing-Jong, LIU Shang-Bin*, SHEN Wenjie*
*Dalian Institute of Chemical Physics, Chinese Academy of Sciences
 Institute of Atomic and Molecular Sciences, "Academia Sinica", Taiwan
 Taiwan University*


Selective adsorption of pyridine on the acidic sites in the 12-MR pores caused dimethyl ether carbonylation to methyl acetate to occur preferentially on the acidic sites in the 8-MR pores of HMOR, which significantly improved the stability of the catalyst.



Chin. J. Catal., 2010, 31: 739–746 doi: 10.1016/S1872-2067(09)60082-6

A Theoretical Investigation of the Structural Properties of Chemically Modified Mo-S-I Nanowires

WEN Shuhao, HOU Zhufeng*, LIU Jianyong*, HE Guozhong
Dalian Institute of Chemical Physics, Chinese Academy of Sciences; Fudan University




The first principle density functional theory calculation results show that the chemical modification of MoSI nanowire will obtain good structural and electronic properties, which provide the basis for chemical modification of the MoSI nanowire materials applying in catalysis.

Chin. J. Catal., 2010, 31: 747–750 doi: 10.1016/S1872-2067(09)60083-8

Synthesis, Characterization, and Performance of $\text{LaZn}_x\text{Fe}_{1-x}\text{O}_3$ Perovskite Nanocatalysts for the Combustion of Toluene

Seyed Ali HOSSEINI*, Mohammad Taghi SADEGHI, Abdolali ALEMI, Aligholi NIAEI, Dariush SALARI, Leila KAFI-AHMADI
University of Tabriz, Iran




We report the synthesis and characterization of nanocrystalline $\text{LaZn}_x\text{Fe}_{1-x}\text{O}_3$ ($x = 0.01, 0.05, 0.1, 0.2,$ and 0.3) perovskites prepared by sol-gel combustion. Their activity for the catalytic combustion of toluene was investigated.

Degradation of Rhodamine B by Fe-Carbon Nanotubes/TiO₂ Composites under UV Light in Aerated Solution


ZHANG Kan, MENG Zeda, OH Wonchun*
Hanseo University, Korea

The responsibilities for the obvious increase of photocatalytic activity indicated that the photoactivity not only benefits from CNT introduced, but also relates to the cooperative effect of the Fe particles via photo-Fenton reaction.



Vapor-Phase Alkylation of Phenol with *Tert*-butyl Alcohol Catalyzed by H₃PO₄/MCM-41



Mehran GHIACI*, Behzad AGHABARARI
Isfahan University of Technology, Iran



The catalytic performance of Al-MCM-41 containing 5, 10, 15, 20, 30, and 35 wt% H₃PO₄ was studied for the vapor phase alkylation of phenol with *tert*-butyl alcohol from 383–493 K.

Hydrated Alkali Metal Halides as Efficient Catalysts for the Synthesis of Cyclic Carbonates from CO₂ and Epoxides


ZHOU Xi, ZHANG Yi, YANG Xiangui, YAO Jie, WANG Gongying*
Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences



Hydrated alkali metal halides show much higher activity than anhydrous alkali metal halides for the reaction between CO₂ and epoxides in the absence of a co-catalyst and a solvent under mild reaction conditions.

Cu-Zn/Al₂O₃ Catalyst for the Hydrogenation of Esters to Alcohols


YUAN Peng, LIU Zhongyi, ZHANG Wanqing, SUN Haijie, LIU Shouchang*
Zhengzhou University



The yield of high alcohols reached above 86.3% over the Cu-Zn/Al₂O₃ catalyst under relatively mild conditions of 240 °C and 10 MPa hydrogen pressure.

Low-Temperature Synthesis and Characterization of AIPO-Cristobalite in Ionic Liquid


XU Renshun, ZHANG Weiping*, HAN Xiuwen, BAO Xinhe*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences



A novel method for the synthesis of AIPO-cristobalite is reported. AIPO-cristobalite can be synthesized at 80 °C for 30 min in the [bmim]Br ionic liquid. The resultant sample is α -AIPO-cristobalite, and most of the Al and P species are connected by Al–O–P tetrahedrons.

Preparation and Characterization of Rh-Au/ γ -Al₂O₃ Three-Way Nanocatalysts

LIU Licheng, ZI Xuehong, DAI Hongxing, ZHAO Zhen, WANG Xiping, HE Hong*
Beijing University of Technology; China University of Petroleum; Dalian University of Technology



A new method, ultrasound-assisted membrane reduction (UAMR), was used to prepare Rh-Au/ γ -Al₂O₃ catalysts. The catalysts exhibited small particles less than 3 nm and high activity in three-way model reactions.


Static Synthesis and Crystallization Mechanism of ZSM-35 Zeolite

JU Xiaohua, FAN Fengtao, TIAN Fuping*, FENG Zhaochi*

Dalian University of Technology;

Dalian Institute of Chemical Physics, Chinese Academy of Sciences


The building units in the precursors were identified as five-membered and six-membered silicate ring species. The aggregation of these ring species yielded ZSM-35 zeolite through the hydrothermal treatment.



Effect of ZrO₂ Addition on the Properties of CuMn₂/Al-Ti Monolithic Catalysts for Benzene Catalytic Combustion

ZHANG Zhiqiang, HE Zhanfeng, WANG Juanyun, WANG Huanyi, LI Jihua, JIANG Yi*, CHEN Junhe, ZHANG Xiaoxia

Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences




After the addition of ZrO₂, the specific surface area, components dispersion, redox ability, and catalytic activity for benzene combustion of CuMn₂Zr_{1.25}-O_x/Al_{0.82}Ti_{0.18}O_x monolithic catalyst increased.

Preparation of (Fe, Ni)-Codoped ZnO and Its Photocatalytic Activity for Degradation of Methyl Orange

FU Tianhua, GAO Qianqian, LIU Fei, DAI Huajun, KOU Xingming*


Sichuan University



The (Fe, Ni)-codoped ZnO photocatalyst was prepared, and its photocatalytic activity for degradation of methyl orange was discussed. (Fe, Ni) co-doping greatly improves the photocatalytic activity of ZnO.

Synthesis of UZM-5 Zeolite by a Hydrothermal Method Using Tetraethylammonium Hydroxide-Tetramethylammonium Chloride as Organic Templates

LIU Yan, LI Yingxia*, LI Jianwei, CHEN Biaohua
Beijing University of Chemical Technology




The synthesis of UZM-5 zeolite by the hydrothermal method in the tetraethylammonium hydroxide (TEAOH)-tetramethyl-ammonium chloride (TMACl) system was studied and mainly focused on the effects of TEAOH, TMACl, and sodium ions on the crystallization process of UZM-5 zeolite.

Effects of Cr Promoter on Structure and Fischer-Tropsch Synthesis Performance over Fe-Based Catalysts


WANG Hulin, YANG Yong*, WANG Hong, QING Ming, XIANG Hongwei, LI Yongwang
Institute of Coal Chemistry, Chinese Academy of Sciences

Cr atoms were incorporated into the lattice of α -Fe₂O₃ and located at the substitutional sites of Fe atoms. With the increase of Cr content in the catalysts, the Fe-rich phase α -(Fe_{1-x}Cr_x)₂O₃ and Cr-rich phase α -(Cr_{1-y}Fe_y)₂O₃ appeared.



Effect of Combination Way of Dielectric Barrier Discharge and CuZSM-5 on NO_x Removal

CHEN Gang, SUN Qi*, SHI Lei, NIU Jinhai, SONG Zhimin
Liaoning Normal University; Dalian University of Technology




The stable species and some short-lived active species formed from the plasma-induced or plasma-catalytic process may be responsible for the synergistic effect between the dielectric barrier discharge (DBD) plasmas and CuZSM-5 catalyst for NO_x removal.

Solid Sulfonic Acid-Promoted Oxidation of Benzyl Alcohol Catalyzed by Isoamyl Nitrite

SHENG Xuebin, MA Hong, LI Decai, HE Jing, XU Jie*

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; PetroChina Urumuqi Petrochemical Company Fibre Mill




The oxidation of benzyl alcohol catalyzed by isoamyl nitrite was promoted by solid sulfonic acid. A significant promotional effect was observed using a solid acid prepared by treating Amberlyst 15 at 300 °C.

Knoevenagel Condensation Catalyzed by Immobilized Ionic Liquid-Proline on SBA-15

SHEN Jiachun, GUO Jianping*, SUN Yanmei, TANG Binyan, CHEN Xiaohua, YIN Dulin*

Hunan Normal University




The IL-Pro/SBA-15 catalyst was prepared using SBA-15 as the support to immobilize ionic liquid-proline (IL-Pro), which was synthesized from L-proline. The catalytic performance of the catalyst was measured for Knoevenagel reaction of benzaldehyde with malononitrile.

Preparation of Ultrafine Y Zeolite from Spent Fluid Catalytic Cracking Catalyst Powders


LIU Xinmei*, LIANG Haining, LI Liang, YANG Tingting, YAN Zifeng
China University of Petroleum

Using the spent Fluid catalytic cracking catalyst powders, the ultrafine Y zeolite was successfully prepared by a hydrothermal crystallization method. The zeolite possessed higher relative crystallinity and higher surface area.



Preparation of Au/SBA-15 and Its Catalytic Activity for CO Oxidation

SU Jixin*, ZHANG Shenping, MA Liyuan, QU Wen, ZHANG Mingbo
Shandong University




Au/SBA-15 catalyst with high activity and anti-sintering property was prepared by a facile deposition-precipitation method, using ammonia water as precipitant which can react with $HAuCl_4$ to form positive charged gold species.

Hydrogenation of Aromatic Ketones to Aromatic Hydrocarbons over SiO_2 -Supported Chitosan Schiff-Base Palladium Catalyst


HE Haifeng, GONG Shuwen*, LIU Lijun, CUI Qingxin, YIN Handong
Liaocheng University

Aromatic aldehydes or ketones can be effectively converted to their corresponding hydrocarbons by catalytic hydrogenolysis at low temperature and under $p(H_2) = 0.1$ MPa using silica-supported chitosan (CS) Schiff-base (SB) palladium SiO_2 -CS-SB-Pd as the catalyst.



Preparation of Chlorine-Free Cu/AC Catalyst and Its Catalytic Properties for Vapor Phase Oxidative Carbonylation of Methanol

WANG Ruiyu, LI Zhong*, ZHENG Huayan, XIE Kechang
Taiyuan University of Technology



When heating $Cu_2(NO_3)(OH)_3/AC$ under N_2 atmosphere, the copper species on the activated carbon surface changed to CuO at $200^\circ C$, to Cu_2O at $300\sim 400^\circ C$, and to Cu^0 at $450^\circ C$. For the oxidative carbonylation of methanol to dimethyl carbonate, the catalytic activity increased in the order of $CuO/AC < Cu_2O/AC < Cu^0/AC$.