

http://www.chxb.cn
ISSN 0253-9837
CN 21-1195/O6
CODEN THHPD3

催化学报

CHINESE JOURNAL OF CATALYSIS

主编 林励吾

Editor-in-Chief LIN Liwu

2012

Vol. 33 No. 5



ISSN 0253-9837



万方数据

中国化学会催化学会会刊
Transaction of the Catalysis Society of China

Science Press



中国科学院科学出版基金资助出版

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

月刊 SCI 收录 2012 年 5 月 第 33 卷 第 5 期



目次

综 述

745 (国际版/封面文章)

二氧化碳作为羧化试剂用于羧酸及其衍生物的合成
张文珍, 吕小兵

757

负载型 Heck 反应催化剂的研究进展
刘鸿飞, 贾志刚, 季生福

研究快讯

768

振荡天平结合微反研究程序升温条件下 SAPO-34 催化的
甲醇转化
袁翠岭, 魏迎旭, 许磊, 李金哲, 徐舒涛, 周游, 陈景润,
王全义, 刘中民

研究论文

771 (国际版)

Bi-MCM-41 催化对氯甲苯选择氧化
赵俊理, 钱广, 李风云, 朱杰, 季生福, 李蕾

777 (国际版)

石墨烯-二氧化钛复合催化剂对光催化性能的提高
赵慧敏, 苏芳, 范新飞, 于洪涛, 吴丹, 全燮

783 (国际版)

低温等离子体协助 $B_2O_3/\gamma-Al_2O_3$ 选择催化还原 NO
于琴琴, 刘彤, 王卉, 肖丽萍, 陈敏, 蒋晓原, 郑小明

790 (国际版)

Ni_2P/SiO_2 和 Ni/SiO_2 催化剂甘油水解反应性能比较: 催
化剂活性及产物选择性影响因素的探讨
黄金花, 陈吉祥

797 (国际版)

Cu 部分取代 $LaNiO_3$ 中的 Ni 对其甲烷干气重整反应性能
的影响
G. R. MORADI, F. KHOSRAVIAN, M. RAHMANZADEH

802 (国际版)

硫化亚铜/四针状氧化锌晶须纳米复合材料的制备及其光
催化性能
吴德智, 范希梅, 代佳, 刘花蓉, 刘红, 张冯章

808

Li 助剂对 Co/AC 催化剂上 CO 加氢制直链混合伯醇反应
性能的影响
裴彦鹏, 丁云杰, 臧娟, 宋宪根, 董文达, 朱何俊, 王涛,
陈维苗

813

V_2O_5/CeO_2 催化剂上甲醇氧化一步法合成二甲氧基甲烷
郭荷芹, 李德宝, 陈从标, 范志宏, 孙子罕

819

具有低 SO_2 氧化活性的 $Pt/Zr,Ti_{1-x}O_2$ 柴油车氧化催化剂

的制备及性能

杨铮铮, 陈永东, 赵明, 周菊发, 龚茂初, 陈耀强

827

$SO_4^{2-}/SnO_2/SBA-15$ 固体酸催化剂的制备及其在环酮缩合
反应中的应用

杨志旺, 洪伟, 王佳, 陈丽娜, 贾娜, 赵磊, 马恒昌,
雷自强

833

Au/MOF 催化剂的制备、表征及其催化三组分偶联反应
刘丽丽, 张鑫, 高金森, 徐春明

842

低温水煤气变换催化剂 Cu/ZrO_2 的制备、表征与性能
阮春晓, 陈崇启, 张燕杰, 林性怡, 詹瑛瑛, 郑起

850

CdS/TiO_2 纳米管可见光催化剂的制备、表征及光催化活性
周强, 苑宝玲, 许东兴, 付明来

857

交联酶聚集体与仿生硅化技术结合制备固定化脂肪酶
姜艳军, 王旗, 王温琴, 周丽亚, 高静

863

单相碳化铁的制备及其表面吸附性质
王瑞雪, 吴宝山, 李永旺

870

超细 Fe-V-O 催化剂上甲苯液相氧化制苯甲醛
张贵泉, 张昕, 祁敏, 林涛, 龚婷

878

硅胶负载的亚胺环钼催化剂的制备、表征及催化性能
杨新丽, 张成军, 戴维林, 刘建平, 韦梅生

885

交联聚苯乙烯微球固载的双齿席夫碱型氧钒(IV) 配合物
催化分子氧氧化苯甲醇
代新, 高保娇, 雷海波

891

多壁碳纳米管固载金鸡纳生物碱季铵盐类手性相转移催
化剂的制备及其催化烷基化反应性能
刘健, 刘奕, 石鑫, 杨启华

898

杂原子 Sn- β 分子筛的脱铝补位两步法制备、表征及其催
化环己酮 Baeyer-Villiger 氧化性能
康自华, 刘海鸥, 张雄福

相关信息

812 第十六届全国催化学术会议(第一轮通知)

905 作者索引



Supported by the Science Publication
Foundation of the CAS

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

Monthly Vol. 33 No. 5 May 2012



Contents


Reviews

Chin. J. Catal., 2012, 33: 745–756 doi: 10.1016/S1872-2067(11)60390-2

Synthesis of Carboxylic Acids and Derivatives Using CO₂ as Carboxylative Reagent

ZHANG Wenzhen*, LÜ Xiaobing
Dalian University of Technology

This review summarized transition metal-catalyzed reactions of CO₂ with carbon nucleophiles, C–H bond and unsaturated organic compounds for the synthesis of carboxylic acid and derivatives using CO₂ as a carboxylative reagent.




Chin. J. Catal., 2012, 33: 757–767 doi: 10.3724/SP.J.1088.2012.20145

Research Progress in Supported Catalysts for Heck Reaction

LIU Hongfei, JIA Zhigang, JI Shengfu*
Beijing University of Chemical Technology

The research progress of the supported Pd-based and non-Pd-based Heck reaction catalysts, which exhibited high catalytic activity and excellent reusable performance, was reviewed.




Communication

Chin. J. Catal., 2012, 33: 768–770 doi: 10.3724/SP.J.1088.2012.20406

Temperature-Programmed Methanol Conversion on a Microscale Setup Equipped with Tapered Element Oscillating Microbalance

YUAN Cuiyu, WEI Yingxu, XU Lei, LI Jinzhe, XU Shutao, ZHOU You, CHEN Jingrun, WANG Quanyi, LIU Zhongmin*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences

Temperature-programmed methanol conversion from 250 to 450 °C goes through the induction period, the deactivation period, and the reactivity recovery period, which results from coke evolution in the SAPO-34 catalyst phase.




Articles

Chin. J. Catal., 2012, 33: 771–776 doi: 10.1016/S1872-2067(11)60367-7

Catalytic Selective Oxidation of 4-Chlorotoluene by Bi-MCM-41

ZHAO Junli, QIAN Guang^{*}, LI Fengyun, ZHU Jie, JI Shengfu, LI Lei
Changzhou University; Jiaxing University; Beijing University of Chemical Technology


A series of bismuth incorporated MCM-41 samples were synthesized. Bi-MCM-41 catalyzed the selective oxidation of 4-chlorotoluene efficiently even on a large scale.



Chin. J. Catal., 2012, 33: 777–782 doi: 10.1016/S1872-2067(11)60374-4

Graphene-TiO₂ Composite Photocatalyst with Enhanced Photocatalytic Performance

ZHAO Huimin^{*}, SU Fang, FAN Xinfei, YU Hongtao, WU Dan, QUAN Xie
Dalian University of Technology




The chemically bonded graphene-TiO₂ nanocomposite photocatalyst was a more stable and better photocatalyst than pure TiO₂. The holes governed the photocatalytic process and the photocatalytic performance of this photocatalyst was improved at high pH value.

Chin. J. Catal., 2012, 33: 783–789 doi: 10.1016/S1872-2067(11)60362-8

Cold Plasma-Assisted Selective Catalytic Reduction of NO over B₂O₃/ γ -Al₂O₃

YU Qinqin, LIU Tong, WANG Hui, XIAO Liping, CHEN Min, JIANG Xiaoyuan, ZHENG Xiaoming^{*}
Zhejiang University




Plasma gas-phase reactions generated NO₂, HCHO, CH₃NO, and CH₃NO₂ as active chemicals. Further reactions of these chemicals on the post-plasma B₂O₃/ γ -Al₂O₃ result in the conversion of NO.

Comparison of Ni₂P/SiO₂ and Ni/SiO₂ for Hydrogenolysis of Glycerol: A Consideration of Factors Influencing Catalyst Activity and Product Selectivity

HUANG Jinhua, CHEN Jixiang*

Tianjin University of Chemical Engineering and Technology




Glycerol was mainly converted to 1,2-propanediol and 1-propanol over Ni₂P/SiO₂, while it was mainly converted to 1,2-propanediol, ethylene glycol, ethanol, and methane over Ni/SiO₂.

Effect of Partial Substitution of Ni by Cu in LaNiO₃ Perovskite Catalyst for Dry Methane Reforming

G. R. MORADI*, F. KHOSRAVIAN, M. RAHMANZADEH

Razi University, Iran


The effect of Cu addition to LaNiO₃ as LaNi_{1-x}Cu_xO₃ (x = 0.2, 0.4, 0.6, 0.8, and 1.0) perovskite type oxides on the dry reforming of methane has been investigated. LaNi_{0.8}Cu_{0.2}O₃ showed the best performance with superior reduction of the nickel.



Preparation and Photocatalytic Properties of Cu₂S/Tetrapod-Like ZnO Whisker Nanocomposites

WU Dezhi, FAN Ximei*, DAI Jia, LIU Huarong, LIU Hong, ZHANG Fengzhang


Southwest Jiaotong University



Coupled Cu₂S/tetrapod-like ZnO whisker (T-ZnOw) catalysts were synthesized by the polyol process. They have a higher activity for the photocatalytic degradation of methyl orange than T-ZnOw.

Effects of Li Promoter on the Performance of Co/AC Catalysts for CO Hydrogenation to Mixed Linear α -Alcohols

PEI Yanpeng, DING Yunjie*, ZANG Juan, SONG Xianguan, DONG Wenda, ZHU Hejun, WANG Tao, CHEN Weimiao
Dalian Institute of Chemical Physics, Chinese Academy of Sciences




The Li additive on the activated carbon (AC) supported Co-based catalysts favored the formation of Co₂C species, which improved the selectivities for mixed linear α -alcohols in CO hydrogenation reaction.

One-Step Oxidation of Methanol to Dimethoxymethane on V₂O₅/CeO₂ Catalyst

GUO Heqin, LI Debao*, CHEN Congbiao, FAN Zhihong, SUN Yuhan*
Institute of Coal Chemistry, Chinese Academy of Sciences;
Shanghai Advanced Research Institute, Chinese Academy of Sciences


The V₂O₅ content influenced the physico-chemical properties of the V₂O₅/CeO₂ catalyst, which strongly affected catalytic performance. As the V₂O₅ content reached 15%, the 15% V₂O₅/CeO₂ catalyst exhibited higher activity for oxidation of methanol to dimethoxymethane.



Preparation and Properties of Pt/Zr_xTi_{1-x}O₂ Catalysts with Low-Level SO₂ Oxidation Activity for Diesel Oxidation

YANG Zhengzheng, CHEN Yongdong*, ZHAO Ming, ZHOU Jufa, GONG Maochu, CHEN Yaoqiang*
Sichuan University, Southwest Petroleum University


The Pt/Zr_{0.1}Ti_{0.9} catalyst exhibits higher catalytic activity for oxidation of CO and C₂H₄, as well as low-level activity for oxidation of SO₂.



Preparation and Application of $\text{SO}_4^{2-}/\text{SnO}_2/\text{SBA-15}$ Solid Acid Catalyst for Acetalization of Cyclic Ketones


YANG Zhiwang*, HONG Wei, WANG Jia, CHEN Lina, JIA Na, ZHAO Lei, MA Hengchang, LEI Ziqiang
Northwest Normal University

$\text{SO}_4^{2-}/\text{SnO}_2/\text{SBA-15}$ was used as an excellent catalyst for acetalization of several cyclic ketones with diols. The catalyst can be regenerated by simple calcination and recycled several times without any significant loss in catalytic activity.



Preparation and Characterization of Metal–Organic Framework Supported Gold Catalysts and Their Catalytic Performance for Three-Component Coupling Reaction


LIU Lili, ZHANG Xin*, GAO Jinsen, XU Chunming*
China University of Petroleum



Gold-metal organic framework (IRMOF-3-SI-Au) bridges the gap between homogeneous and heterogeneous gold catalysts, and the catalyst is highly active and selective for three-component coupling reaction.

Cu/ZrO₂ Catalyst for Low-Temperature Water-Gas Shift Reaction: Preparation, Characterization and Performance

RUAN Chunxiao, CHEN Chongqi, ZHANG Yanjie, LIN Xingyi, ZHAN Yingying, ZHENG Qi*
Fuzhou University




A series of Cu/ZrO_2 catalyst samples were fabricated by hydrothermal synthesis combined with a deposition-precipitation method. The effect of preparation parameters on the catalytic performance of these catalyst samples was investigated.

Synthesis, Characterization and Photocatalytic Performance of CdS/TiO₂ Nanotube Photocatalyst


ZHOU Qiang, YUAN Baoling*, XU Dongxing, FU Minglai
Huaqiao University; Fuzhou University;
Institute of Urban Environment, Chinese Academy of Sciences

CdS quantum dots have been successfully deposited onto the surfaces of TiO₂ nanotubes (TNTs) to form high-active CdS/TNTs photocatalyst by use of a simple bifunctional organic linker, thiolactic acid.



Preparation of Immobilized Lipase through Combination of CLEAs and Biomimetic Silicification


JIANG Yanjun, WANG Qi, WANG Wenqin, ZHOU Liya, GAO Jing*
Hebei University of Technology; Institute of Process Engineering, Chinese Academy of Sciences



The hybrid biocatalyst was prepared by the combination of cross-linked enzyme aggregates (CLEAs) and biomimetic silicification, which may provide a simple technique for enzyme immobilization.

Synthesis of Single-Phase Iron Carbides and Their Adsorption Performance


WANG Ruixue, WU Baoshan*, LI Yongwang
Institute of Coal Chemistry, Chinese Academy of Sciences



Iron carbides with single-phases, including ϵ -Fe₂C, γ -Fe₅C₂, and θ -Fe₃C, were synthesized in different carbonization gases at different temperatures. The effects of carbonization conditions on the structural properties, carbon deposition and adsorption properties of the iron carbides were studied.

Superfine Fe-V-O Catalyst for Liquid-Phase Oxidation of Toluene to Benzaldehyde

ZHANG Guiquan, ZHANG Xin*, QI Min, LIN Tao, GONG Ting
Northwest University




Hydrothermally synthesized Fe-V-O showed the highest benzaldehyde yield and better reusability in liquid-phase oxidation of toluene by H₂O₂ to benzaldehyde due to its smaller particle size, higher V/Fe surface ratio, and moderate reducibility.

Preparation and Characterization of Silica-Supported Imine Palladacycle Complex and Its Catalytic Performance


YANG Xinli*, ZHANG Chengjun, DAI Weilin*, LIU Jianping,
WEI Meisheng
Henan University of Technology; Fudan University

The silica-supported imine palladacycle complex was synthesized. The complex was a promising novel catalyst in the Heck reaction of iodobenzene with acrylamide to *trans*-products.



Catalytic Oxidation of Benzyl Alcohol by Dioxygen over Bidentate Schiff Base-Type Oxovanadium(IV) Complex Immobilized on Crosslinked Polystyrene Microspheres

DAI Xin, GAO Baojiao*, LEI Haibo
North University of China; Weifang Institute of Supervision & Inspection on Product Quality




An immobilized Schiff base-type oxovanadium(IV) complex, CPS-[VO(SAAM)₂], was prepared. In benzyl alcohol oxidation by dioxygen, the heterogeneous oxovanadium(IV) complex catalyst has high catalytic activity and excellent catalytic selectivity.

Immobilization of Cinchona Quaternary Ammonium Salts as the Chiral Phase Transfer Catalysts on Multi-walled Carbon Nanotubes and Their Application in Enantioselective Alkylation

LIU Jian, LIU Yan*, SHI Xin*, YANG Qihua*

Liaoning Normal University; Dalian Institute of Chemical Physics, Chinese Academy of Sciences




A new catalyst CPTC/MWCNTs was synthesized by adsorption of pyrene-tagged cinchona quaternary ammonium salt (CPTC) onto multi-walled carbon nanotubes (MWCNTs) via π - π stacking interaction. CPTC/MWCNTs was successfully employed in asymmetric alkylation of different halohydrocarbons with *N*-(diphenylmethene)glycine *tert*-butyl ester.

Preparation and Characterization of Sn- β Zeolites by a Two-Step Postsynthesis Method and Their Catalytic Performance for Baeyer-Villiger Oxidation of Cyclohexanone

KANG Zihua, LIU Hai'ou, ZHANG Xiongfu*

Dalian University of Technology



Sn- β zeolites were prepared by a two-step postsynthesis method including first creating vacant T-sites by dealumination of Al- β zeolites and then inserting Sn^{4+} into the T-sites of the β zeolites by impregnation and calcination. They show impressive catalytic performance in Baeyer-Villiger oxidation of cyclohexanone.