

http://www.chxb.cn
ISSN 0253-9837
CN 21-1195/O6
CODEN THHPD3

催化学报


CHINESE JOURNAL OF CATALYSIS

主编 林励吾

Editor-in-Chief LIN Liwu

2012

Vol. 33 No. 7



ISSN 0253-9837



中国化学会催化学会会刊
Transaction of the Catalysis Society of China


Science Press

万方数据



中国科学院科学出版基金资助出版

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

月刊 SCI 收录 2012年7月 第33卷 第7期



目次

研究快讯

1069 (国际版/封面文章)

氧化铈的结构对其热稳定性及催化丙烷氧化脱氢反应性能的影响

高旭锋, 湛春林, 任士远, 张建, 苏党生

1075 (国际版)

柠檬酸络合法制备 Ba 促进 ZrO₂ 负载 Ru 催化剂上氨合成反应性能

林建新, 王自庆, 张留明, 倪军, 王榕, 魏可镁

1080 (国际版)

水蒸气处理对 Ni₂P/SiO₂ 催化剂催化氯苯加氢脱氯反应的促进作用

郭提, 陈吉祥, 李克伦

研究论文

1086 (国际版)

CeAlPO-5 分子筛催化剂上二苯基甲烷选择性氧化生成二苯甲酮

Subbiah DEVIKA, Muthiahpillai PALANICHAMY, Velayutham MURUGESAN

1095 (国际版)

硅胶键合 *N*-丙基二乙烯氨基磺酸作为可回收固体酸催化剂催化合成 α -氨基腈

Tahere RAHI, Mojtaba BAGHERNEJAD, Khodabakhsh NIKNAM

1101 (国际版)

可分散的 In₂O₃/Ta₂O₅ 复合光催化剂的制备及其光催化制氢性能

许蕾蕾, 倪磊, 施伟东, 官建国

1109 (国际版)

Sr 取代 LaFeO₃ 钙钛矿的结构性质和催化性能

张晓静, 李华举, 李勇, 申文杰

1115 (国际版)

炼钢渣和炭的混合物用于微波辅助甲烷干气重整反应

Jose M. BERMUDEZ, Beatriz FIDALGO, Ana ARENILLAS, J. Angel MENENDEZ

1119 (国际版)

费托合成反应中纳米铁催化剂上 La 和 Ba 助剂的协同效应

Yahya ZAMANI, Mehdi BAKAVOLI, Mohamad RAHIMIZADEH, Ali MOHAJERI, Seyed Mohamad SEYEDI

1125 (国际版)

Au/CeO₂/SiO₂ 催化 CO 低温氧化反应过程中 CeO₂ 的作用

张慧丽, 任丽会, 陆安慧, 李文翠

1133 (国际版)

助剂对 L-脯氨酸催化直接不对称 Aldol 反应的影响

骆建轻, 谭蓉, 孔瑜, 黎成勇, 银董红

1139 (国际版)

碱土金属氧化物对丙三醇和苯胺气相合成 3-甲基吡啶的 Cu/SiO₂-Al₂O₃ 催化剂的作用

王兆宇, 李晓辉, 张跃, 石雷, 孙琪

1146 (国际版)

聚乙二醇 400-水介质中水溶性钌膦二胺催化苯叉丙酮的不对称加氢反应

秦瑞香, 王金波, 熊伟, 冯建, 刘德蓉, 陈华

1154 (国际版)

(1*S*,2*S*)-1,2-二苯基乙二胺修饰 Ir/SiO₂ 催化苯乙酮及其衍生物不对称加氢

杨朝芬, 杨俊, 孙晓东, 朱艳琴, 王齐, 陈华

1161 (国际版)

离子液体对 β -糖苷酶催化合成红景天苷的影响

毕艳红, 王朝宇, 茅燕勇, 郑尚永, 张海江, 时号

1166 (国际版)

颗粒内传质参数作为工具用于催化剂颗粒的设计

L. PETROV, M. DAOUS, Y. ALHAMED, A. AL-ZAHRANI, Kh. MAXIMOV

1176

镁铝混合氧化物负载镍催化剂上液化石油气的预重整

王新星, 汪学广, 尚兴付, 聂望欣, 邹秀晶, 鲁雄钢, 丁伟中

1183

Rh/SiO₂ 催化剂上甲烷部分氧化制合成气的反应机理

温在恭, 李虎, 翁维正, 夏文生, 黄传放, 万惠霖

1191

硝酸水热处理活性炭对其负载的 Ba-Ru-K 氨合成催化剂性能的影响

冯国全, 蓝国钧, 李琰, 韩文锋, 刘化章

1198

4-*N,N*-二甲基吡啶促进乙酸钴-溴苯催化甲苯氧化

张展, 高进, 马红, 徐杰

1203

草酸处理对丙烷氨氧化催化剂 Sb-V-O 结构和催化性能的影响

汪国军, 郭耘, 卢冠志

1209

Ag@AgBr 光催化剂的制备及其可见光催化降解亚甲基蓝反应性能

聂龙辉, 黄征青, 徐洪涛, 张旺喜, 杨柏蕊, 方磊, 李帅华

1217

超细 NaY 分子筛的深度脱铝

王希龙, 宋金娜, 叶修群, 顾海芳, 黄曜, 牛国兴

1224

酸性聚乙烯基吡咯烷酮-杂多酸杂化催化剂的合成及其催化酯化反应性能

冷炎, 仇学谦, 蒋平平, 王军

1229

Ni-Cu/Al₂O₃ 催化剂上顺酐液相选择加氢制丁二酸酐反应性能

王达, 张因, 李海涛, 赵丽丽, 张鸿喜, 赵永祥

1236

干胶法制备钛硅沸石及其催化性能

王维海, 李钢, 刘丽萍, 陈永英

相关信息

1145 《催化学报》2011 年 SCI 影响因子首次突破 1.0

1242 作者索引

www.chxb.cn



Supported by the Science Publication
Foundation of the CAS

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

Monthly Vol. 33 No. 7 July 2012



Contents

Communications


Chin. J. Catal., 2012, 33: 1069–1074 doi: 10.1016/S1872-2067(11)60404-X

Structural Effects of Cerium Oxides on Their Thermal Stability and Catalytic Performance in Propane Oxidation Dehydrogenation

GAO Xufeng, CHEN Chunlin, REN Shiyuan, ZHANG Jian*,
SU Dangsheng*

Institute of Metal Research, Chinese Academy of Sciences

This communication presents the change of cerium oxides nanostructures with the thermal treatment, being important to make clear the relationship between structure and thermal stability and catalytic performance.




Chin. J. Catal., 2012, 33: 1075–1079 doi: 10.1016/S1872-2067(11)60413-0

Ammonia Synthesis over Ruthenium Catalysts using Barium-Doped Zirconia as Supports Prepared by Citric Acid Method

LIN Jianxin*, WANG Ziqing, ZHANG Liuming, NI Jun,
WANG Rong, WEI Kemei

Fuzhou University

Ru/Ba-ZrO₂ was an excellent catalyst for ammonia synthesis especially under mild conditions. This was attributed to the presence of BaZrO₃ with high basicity and ability to conduct electrons.




Chin. J. Catal., 2012, 33: 1080–1085 doi: 10.1016/S1872-2067(11)60418-X

Promotion Effect of Steam Treatment on Activity of Ni₂P/SiO₂ for Hydrodechlorination of Chlorobenzene

GUO Ti, CHEN Jixiang*, LI Kelun

Tianjin University




A promoting effect of steam treatment on the activity of Ni₂P/SiO₂ for hydrodechlorination of chlorobenzene was found. This is probably related to synergism between Ni sites and P–OH groups.

Chin. J. Catal., 2012, 33: 1086–1094 doi: 10.1016/S1872-2067(11)60401-4

Selective Oxidation of Diphenylmethane to Benzophenone over CeAlPO-5 Molecular Sieves

Subbiah DEVIKA, Muthiahpillai PALANICHAMY,
Velayutham MURUGESAN*
Anna University, India


Selective oxidation of diphenylmethane was carried over CeAlPO-5 (Al/Ce = 25, 50, 75, 100, and 125) molecular sieves between 250 and 325 °C. The main product of oxidation was benzophenone. The catalytic activity increased with decrease in the Al/Ce ratio of the catalyst.



Chin. J. Catal., 2012, 33: 1095–1100 doi: 10.1016/S1872-2067(11)60383-5

Silica-Bonded *N*-Propyldiethylenetriamine Sulfamic Acid as a Recyclable Solid Acid Catalyst for the Synthesis of α -Aminonitriles

Tahere RAHI, Mojtaba BAGHERNEJAD, Khodabakhsh NIKNAM*
Gachsaran Branch, Islamic Azad University, Iran; Persian Gulf University, Iran




α -Aminonitriles were synthesized using silica-bonded *N*-propyldiethylenetriamine sulfamic acid as a recyclable solid acid catalyst for condensation of an aldehyde, amine, and trimethylsilyl cyanide at room temperature.

Chin. J. Catal., 2012, 33: 1101–1108 doi: 10.1016/S1872-2067(11)60382-3

Photocatalytic Activity for Hydrogen Evolution over Well-Dispersed Heterostructure In₂O₃/Ta₂O₅ Composites


XU Leilei, NI Lei, SHI Weidong, GUAN Jianguo*
Wuhan University of Technology

Well-dispersed In₂O₃/Ta₂O₅ composites exhibit enhanced photocatalytic activity for hydrogen production under simulant solar light irradiation because of effective photogenerated charge-carrier separation between In₂O₃ and Ta₂O₅.



Structural Properties and Catalytic Activity of Sr-Substituted LaFeO₃ Perovskite


ZHANG Xiaojing, LI Huaaju, LI Yong, SHEN Wenjie*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences



Sr substitution in LaFeO₃ induced the generation of oxygen vacancies, which facilitated the dissociation of gaseous oxygen and diffusion of lattice oxygen, and enhanced the activity for CO oxidation and methane combustion.

Mixtures of Steel-Making Slag and Carbons as Catalyst for Microwave-Assisted Dry Reforming of CH₄

Jose M. BERMUDEZ, Beatriz FIDALGO, Ana ARENILLAS, J. Angel MENENDEZ*
Instituto Nacional del Carbon, Spain




The use of steel-making slag as catalysts for microwave-assisted dry reforming of CH₄ is investigated. Two carbon materials, mixtures of the carbon materials and Fe-rich slag, and mixtures of the carbon materials and Ni/Al₂O₃ are tested as catalysts.

Synergetic Effect of La and Ba Promoters on Nanostructured Iron Catalyst in Fischer-Tropsch Synthesis

Yahya ZAMANI, Mehdi BAKAVOLI*, Mohamad RAHIMIZADEH,
Ali MOHAJERI, Seyed Mohamad SEYEDI
Ferdowsi University of Mashhad, Iran;
National Iranian Oil Company, Iran


The effect of promoters such as Ba and La has been investigated on the conventional nanostructured iron catalyst in Fischer-Tropsch synthesis (FTS). The results indicate that adding promoters could improve activity of Fe catalysts for FTS and water-gas shift reaction and lower the gas fraction at the outlet. In addition, both Ba- and La-promoted Fe catalysts exhibit the highest activity due to the synergetic effect.



Role of CeO₂ in Three-Component Au/CeO₂/SiO₂ Composite Catalyst for Low-Temperature CO Oxidation

ZHANG Huili, REN Lihui, LU Anhui, LI Wencui*

Dalian University of Technology




Hierarchical composite nanostructures composed of Au, CeO₂, and SiO₂ were fabricated by sequentially depositing ceria nanoparticles through impregnation and calcination and then gold nanoparticles through deposition-precipitation on hierarchical monolithic silica (HMS). The presence of ceria had a significant effect on controlled on target deposition and the stabilization of small metallic gold nanoparticles on the support.

Effect of Additives on L-Proline Catalyzed Direct Asymmetric Aldol Reactions

LUO Jianqing, TAN Rong*, KONG Yu, LI Chengyong, YIN Donghong*

Hunan Normal University; China Tobacco Hunan Industrial Corporation




Cheap catechol promoted the L-proline catalyzed aldol reaction efficiently through hydrogen-bonds interaction, which decreased the amount of L-proline and increased the yield of aldol products.

Effect of Alkaline-Earth Metal Oxides on Cu/SiO₂-Al₂O₃ Catalyst for Vapor-Phase Synthesis of 3-Methylindole from Glycerol and Aniline

WANG Zhaoyu, LI Xiaohui, ZHANG Yue, SHI Lei*, SUN Qi


Liaoning Normal University



Alkaline-earth oxides introduced into a Cu/SiO₂-Al₂O₃ catalyst for the vapor-phase synthesis of 3-methylindole from glycerol and aniline and MgO improved its performance.

Green and Recyclable Medium for Asymmetric Hydrogenation of Benzalacetone Catalyzed by $\text{RuCl}_2(\text{TPPTS})_2$ -(S,S)-DPENDS


QIN Ruixiang, WANG Jinbo*, XIONG Wei, FENG Jian, LIU Derong, CHEN Hua*
Chongqing University of Science and Technology; Sichuan University



PEG400/H₂O was an efficient recyclable reaction medium for asymmetric hydrogenation of benzalacetone catalyzed by $\text{RuCl}_2(\text{TPPTS})_2$ -(S,S)-DPENDS. Utilizing this medium chemoselectivity for 4-phenyl-3-buten-2-ol was 98.5% and ee value of 74.3% was achieved. The catalyst immobilized on PEG400/H₂O could be recycled and used repeatedly without significant loss of chemoselectivity and enantioselectivity.

(1*S*,2*S*)-Diphenylethylenediamine Modified Ir/SiO₂ Catalyst for Asymmetric Hydrogenation of Acetophenone and Its Derivatives


YANG Chaofen, YANG Jun, SUN Xiaodong, ZHU Yanqin, WANG Qi, CHEN Hua*
Kunming University of Science and Technology; Sichuan University



5%Ir/SiO₂ prepared under basic conditions exhibited good catalytic performance in the asymmetric hydrogenation of acetophenone and its derivatives.

Ionic Liquid Effects on the Activity of β -Glycosidase for the Synthesis of Salidroside in Co-solvent Systems

BI Yanhong, WANG Zhaoyu*, MAO Yanyong, ZHENG Shangyong, ZHANG Haijiang, SHI Hao
Huaiyin Institute of Technology




The effects of 1-alkylimidazolium-based ionic liquids possessing different alkyl chain lengths (C₂–C₁₀) and various anions (BF₄⁻, PF₆⁻, Cl⁻, Br⁻, and I⁻) on β -glycosidase activity for the synthesis of salidroside were investigated.

Use of Intraparticle Mass Transfer Parameters as a Design Tool for Catalyst Pellets

L. PETROV*, M. DAOUS, Y. ALHAMED, A. AL-ZAHRANI, Kh. MAXIMOV

King Abdulaziz University, Kingdom of Saudi Arabia; Institute of Catalysis, Bulgarian Academy of Sciences, Bulgaria




The preparation of catalyst pellets with small pore tortuosity is one of the objectives in selecting pelletizing technology and pelletizing conditions. The intra-particle mass transfer parameters are helpful tool for achieving this aim.

Pre-reforming of Liquefied Petroleum Gas over Magnesium Aluminum Mixed Oxide Supported Nickel Catalysts

WANG Xinxing, WANG Xueguang*, SHANG Xingfu, NIE Wangxin, ZOU Xiujing, LU Xionggang, DING Weizhong
Shanghai University


Ni catalysts supported on Mg-Al mixed oxides prepared by a co-precipitation-impregnation method showed excellent catalytic performance and stability for LPG pre-reforming at a low steam/carbon molar ratio of 2.0.



Reaction Mechanism for Partial Oxidation of Methane to Synthesis Gas over Rh/SiO₂ Catalyst


WEN Zaigong, LI Hu, WENG Weizheng*, XIA Wensheng, HUANG Chuanjing, WAN Huilin*
Xiamen University

Pyrolysis of methane on reduced rhodium sites followed by coupling of two surface hydrogen atoms to H₂ and partial oxidation of surface carbon species to CO are the major reactions responsible for synthesis gas formation in the oxidation zone of the Rh/SiO₂ catalyst bed.



Effect of Hydrothermal Treatment of Activated Carbon by Nitric Acid on Activity of Ba-Ru-K/AC Catalyst for Ammonia Synthesis

FENG Guoquan, LAN Guojun, LI Ying, HAN Wenfeng, LIU Huazhang*
Zhejiang University of Technology




The hydrothermal treatment with HNO_3 is firstly used to adjust the amount of surface oxygen functional groups on activated carbon to a desired degree. The degree of functionalization is found to be strongly dependent on the concentration of HNO_3 .

4-*N,N*-Dimethylamino Pyridine Promoted Oxidation of Toluene Catalyzed by Cobalt Acetate and Benzyl Bromide


ZHANG Zhan, GAO Jin, MA Hong, XU Jie*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences

The promoting effect of 4-*N,N*-dimethylamino pyridine (DMAP) to cobalt acetate and benzyl bromide catalyzed oxidation of toluene was due to the in situ formation of quaternary pyridinium salts.



Effect of Oxalic Acid on the Structure and Catalytic Performance of Sb-V-O Catalyst for Propane Ammoxidation


WANG Guojun, GUO Yun, LU Guanzhong*
East China University of Science and Technology; Shanghai Research Institute of Petrochemical Technology



The addition of oxalic acid in the VSb_3 mixed oxide preparation process can keep V at a low valence and generate a higher amount of SbVO_4 phase, which is the key active phase for propane ammoxidation.

Synthesis of Ag@AgBr Photocatalyst and Its Performance for Degradation of Methylene Blue under Visible-Light Irridiation

NIE Longhui*, HUANG Zhengqing, XU Hongtao, ZHANG Wangxi, YANG Borui, FANG Lei, LI Shuaihua
Hubei University of Technology




The photocatalytic activity of Ag@AgBr prepared by the deposition-precipitation and photo-reduction method reached 96% in 10 mg/L methylene blue aqueous solution under the visible-light ($\lambda > 420 \text{ nm}$) irradiation, and it showed good stability in five-cycle test.

Deep Dealumination of Ultrafine NaY Zeolite


WANG Xilong, SONG Jinna, YE Xiuqun, GU Haifang,
HUANG Yao*, NIU Guoxing*
Fudan University

Deeply dealuminated ultrafine Y zeolite was prepared successfully by the combination of dealumination and silicon insertion, showing its high framework Si/Al ratio of 27.3 and surface area of 581.9 m^2/g .



Synthesis of Polyvinyl Pyrrolidone-Heteropolyacid Hybrid Catalyst and Its Catalytic Activity for Esterification Reactions


LENG Yan*, QIU Xueqian, JIANG Pingping, WANG Jun
Jiangnan University; Nanjing University of Technology



A new heteropolyacid-based acidic hybrid synthesized by combining polyvinyl pyrrolidone with heteropolyanion was revealed to be an efficient recyclable solid catalyst for esterification of acetic acid with *n*-butanol.

Selective Hydrogenation of Maleic Anhydride to Succinic Anhydride in Liquid Phase over Ni-Cu/Al₂O₃ Catalyst


WANG Da, ZHANG Yin, LI Haitao, ZHAO Lili, ZHANG Hongxi, ZHAO Yongxiang^{*}
Shanxi University



The introduction of Cu to Ni/Al₂O₃ catalyst effectively improved the activity for C=C bond hydrogenation and inhibited the hydrogenation of C=O bond.

Titanium Silicalite Synthesized by Dry Gel Conversion Method and Its Catalytic Performance

WANG Weihai, LI Gang^{*}, LIU Liping, CHEN Yongying
Dalian University of Technology



TS-1 and hierarchical TS-1 were prepared by dry gel conversion technique in TPABr synthetic system. Hierarchical TS-1 could exhibit higher catalytic performance than TS-1 in both thiophene and benzothiophene oxidation.