

http://www.chxb.cn
ISSN 0253-9837
CN 21-1195/O6
CODEN THHPD3

催化学报


Chinese Journal of Catalysis

主编 林励吾

Editor-in-Chief LIN Liwu

2013

Vol. 34 No. 2




ISSN 0253-9837



9 770253 983139

万方数据

中国化学会催化学会会刊
Transaction of the Catalysis Society of China



中国科学院科学出版基金资助出版

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

月刊 SCI收录 2013年2月 第34卷 第2期



目次

综 述

351 (中)

生物制造不同立体构型2,3-丁二醇: 合成机理与实现方法
沈梦秋, 纪晓俊, 聂志奎, 夏志芳, 杨晗, 黄和

研究论文

283 (英/中/封面文章)

焙烧与预处理条件对 Co_3O_4 催化氧化CO性能的影响
余运波, 赵娇娇, 韩雪, 张燕, 秦秀波, 王宝义

294 (英)

ZSM-5沸石结晶度对乙苯叔丁基化对位选择性的影响
PUSHPARAJ Hemalatha, MANI Ganesh, MUTHIAHPILLAI
Palanichamy, VELAYUTHAM Murugesan, PARK Yong-Ki,
CHOI Won Choon, JANG Hyun Tae

305 (英)

Ni掺杂对纳米结构牡丹花状 CeO_2 材料催化特性的影响
仙存妮, 王少飞, 孙春文, 李泓, 陈晓惠, 陈立泉

313 (英/中)

催化臭氧化降解邻苯二甲酸二甲酯中催化剂构效关系
王建兵, 王灿, 杨春丽, 王国庆, 祝万鹏

322 (英)

镍促进 CuO-CeO_2 催化剂的结构表征及低温CO氧化活性
陈国星, 李巧灵, 魏育才, 方维平, 杨意泉

330 (英)

复合氧化物载体对镍基催化剂上CO甲烷化反应性能的影响
张罕, 董芸芸, 方维平, 连奕新

336 (英)

Au/NTS-1催化丙烯气相直接环氧化
刘义武, 张小明, 索继栓

341 (英/中)

Cu掺杂对介孔 $\text{VO}_x\text{-TiO}_2$ 催化苯羟基化制苯酚的影响
徐丹, 贾丽华, 郭祥峰

361 (中)

碱土金属对锆基钙钛矿材料负载钨催化剂氨合成性能的影响
王自庆, 马运翠, 林建新, 王榕, 魏可镁

367 (中)

液相沉积法制备可磁分离复合光催化剂纳米球及其催化性能
许士洪, 谭东栋, 鲁巍, 时鹏辉, 毕得福, 马春燕, 上官文峰

373 (中)

Cr掺杂对中孔 MgF_2 酸性及孔结构的影响
牛怀成, 李利春, 李瑛, 郭荔, 唐浩东, 韩文锋, 刘化章

379 (中)

介孔Ni- $\beta\text{-Mo}_2\text{C/SBA-16}$ 催化剂在 CH_4/CO_2 重整制合成气反应中的催化性能
璁莫汗, 付晓娟, 雷艳秋, 苏海全

385 (中)

Pt/BiOCl纳米片的制备、表征及其光催化性能
余长林, 陈建钊, 操芳芳, 李鑫, 樊启哲, YU Jimmy C,
魏龙福

391 (中)

Ru-Fe/C催化剂上邻氯硝基苯原位液相加氢性能
许响生, 陈傲昂, 周莉, 李小青, 顾辉子, 严新焕

相关信息

293 2nd International Congress on Catalysis for Biorefineries
(CatBior 2013)

321 第二届国际生物质催化炼制大会(CatBior 2013)第一轮
通知

397 作者索引

英文全文电子版(国际版)由Elsevier出版社在ScienceDirect上出版
<http://www.sciencedirect.com/science/journal/18722067>
<http://www.elsevier.com/locate/chnjc>
<http://www.chxb.cn>



Supported by the Science Publication
Foundation of the CAS

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

Monthly Vol. 34 No. 2 February 2013




Graphical Contents

Review

Chin. J. Catal., 2013, 34: 351–360 doi: 10.3724/SP.J.1088.2013.20737

Biotechnological production of 2,3-butanediol stereoisomers: synthetic mechanism and realized methods

SHEN Mengqiu, JI Xiaojun*, NIE Zhikui, XIA Zhifang, YANG Han, HUANG He*
Nanjing University of Technology




The biological routes for the production of pure 2,3-butanediol stereoisomers, including using the methods of whole cell catalysis and the emerging synthetic biology, was reviewed. In contrast to the conventional chemical methods, the biological methods own their great advantages.

Articles

Chin. J. Catal., 2013, 34: 283–293 doi: 10.1016/S1872-2067(11)60484-1

Influence of Calcination and Pretreatment Conditions on the Activity of Co_3O_4 for CO Oxidation

YU Yunbo*, ZHAO Jiaojiao, HAN Xue, ZHANG Yan, QIN Xiubo, WANG Baoyi
*Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences;
Institute of High Energy Physics, Chinese Academy of Sciences*




Pretreatment of Co_3O_4 in N_2 at moderate temperatures promotes the formation of oxygen vacancy clusters, favoring the adsorption of oxygen molecules and guaranteeing a long durability for CO oxidation.

Effects of crystallinity of ZSM-5 zeolite on *para*-selective *tert*-butylation of ethylbenzene

PUSHPARAJ Hemalatha, MANI Ganesh, MUTHIAHPILLAI Palanichamy, VELAYUTHAM Murugesan, PARK Yong-Ki, CHOI Won Choon, JANG Hyun Tae*

Hanseo University, South Korea; Anna University, India; Korea Research Institute of Chemical Technology, South Korea




A fluoride medium offers defect-free, highly crystalline ZSM-5 crystals. High crystallinity confers high *para* selectivity (> 90%) in *tert*-butylation of ethylbenzene. A fluoride medium is better than an alkaline medium for the commercial production of *para*-selective ZSM-5 catalysts.

Effect of Ni doping on the catalytic properties of nanostructured peony-like CeO₂

XIAN Cunni, WANG Shaofei, SUN Chunwen, LI Hong*, CHAN Suiwai, CHEN Liquan

Institute of Physics, Chinese Academy of Sciences, China; Columbia University, USA




Oxygen vacancies are generated in bulk ceria after Ni doping, which promotes the reducibility of peony-like CeO₂, and hence enhances the catalytic activity for CO oxidation.

Relationship between the structure and activity of ruthenium catalysts in the catalytic ozonation of dimethyl phthalate

WANG Jianbing*, WANG Can, YANG Chunli, WANG Guoqing, ZHU Wanpeng


China University of Mining and Technology, Beijing Campus; Tsinghua University



The surface structure of the activated carbon (AC) support influenced the activity of Ru/AC catalysts in dimethyl phthalate ozonation. Microwave heating during catalyst preparation changed the catalyst activity by a modification of its surface structure.

Low temperature CO oxidation on Ni-promoted CuO-CeO₂ catalysts


CHEN Guoxing, LI Qiaoling, WEI Yucai, FANG Weiping,
YANG Yiquan*
Xiamen University



The high catalytic activity of Ni-promoted CuO-CeO₂ is due to the promoter giving increased amounts of Cu⁺ in the catalyst and the formation of solid solutions of Cu-O-Ce and Ni-O-Ce.

Effects of composite oxide supports on catalytic performance of Ni-based catalysts for CO methanation


ZHANG Han, DONG Yunyun, FANG Weiping*, LIAN Yixin*
Xiamen University



NiO/MO_x-Al₂O₃ (M = Mg, Si, Zr) catalysts for CO methanation, prepared using a modified grinding-mixing method, have higher catalytic activities than that of a conventional NiO/Al₂O₃ catalyst. This is attributed to the weakening of Ni-Al interactions after adding MO_x.

Gold supported on nitrogen-incorporated TS-1 for gas-phase epoxidation of propylene


LIU Yiwu, ZHANG Xiaoming*, SUO Jishuan
Chengdu Institute of Organic Chemistry, Chinese Academy of Sciences; Neijiang Normal University



A novel gold catalyst was prepared by immobilization of gold nanoparticles on nitrogen-incorporated TS-1. This catalyst exhibits an excellent catalytic capacity for gas-phase epoxidation of propylene using H₂ and O₂. Nitrogen-incorporation into TS-1 improved both gold loading and dispersion, and decreased the acidic sites of the support surface.

Cu-doped mesoporous VO_x-TiO₂ for catalytic hydroxylation of benzene to phenol


XU Dan, JIA Lihua*, GUO Xiangfeng*
Qiqihar University



Incorporation of Cu additives into a VO_x/TiO₂ catalyst improved the reducibility of VO_x species, while Cu helped the monodispersion of VO_x species on the TiO₂ support surface.

Effect of alkali earth mentals on performance of zirconium-based perovskite composite oxides supported ruthenium for ammonia synthesis

WANG Ziqing, MA Yuncui, LIN Jianxin*, WANG Rong, WEI Kemei
Fuzhou University




BaZrO₃ was an excellent support for Ru-based catalyst for ammonia synthesis compared with CaZrO₃ and SrZrO₃, which could significantly inhibit the adsorption of H₂ and facilitate the cleavage of N₂.

Photocatalytic properties of magnetically separable composite photocatalyst nanospheres prepared by liquid-phase deposition

XU Shihong*, TAN Dongdong, LU Wei, SHI Penghui, BI Defu, MA Chunyan, SHANGGUAN Wenfeng
Donghua University; Beijing General Municipal Engineering Design & Research Institute; Shanghai Jiao Tong University


Surfactant AOT



A novel photocatalyst nanosphere TiO₂@SiO₂@NiFe₂O₄ was prepared by a reverse micelle method and liquid phase deposition technique. The prepared photocatalyst nanospheres show high photocatalytic activity.

Effect of Cr-doping on the acidity and pore structure of mesoporous magnesium fluoride

NIU Huaicheng, LI Lichun, LI Ying*, GUO Li*, TANG Haodong, HAN Wenfeng, LIU Huazhang
Zhejiang University of Technology; Zhejiang Chemical Industry Research Institute Co., Ltd.




The Cr-doping in mesoporous magnesium fluoride prepared by co-precipitation increases the acidity and the specific surface area of magnesium fluoride and thus increases the catalytic performance in CHClF₂ disproportionation.

Catalytic performance of mesoporous material supported bimetallic carbide Ni-β-Mo₂C/SBA-16 catalyst for CH₄/CO₂ reforming to syngas

Naomohan, FU Xiaojuan, LEI Yanqiu, SU Haiquan*
Inner Mongolia University


The catalyst Ni-β-Mo₂C/SBA-16 in methane/carbon dioxide reforming reaction, which establishes carbonization-oxidation circulation, exhibited high catalytic activity and remarkable anti-coke effect.



Preparation, characterization, and photocatalytic properties of Pt/BiOCl nanoplates

YU Changlin*, CHEN Jianchai, CAO Fangfang, LI Xin, FAN Qizhe, YU Jimmy C, WEI Longfu
Jiangxi University of Science and Technology; Fuzhou University; The Chinese University of Hong Kong


The presence of Pt nanoparticles could effectively separate the photo-generated e⁻/h⁺ pairs and result in the plasmon photocatalysis under visible light irradiation.



Catalytic stability of *ortho*-chloronitrobenzene hydrogenation on Ru-Fe/C catalyst

XU Xiangsheng, CHEN Ao'ang, ZHOU Li, LI Xiaoqing, GU Huizi, YAN Xinhuan*
Zhejiang University of Technology

CO accumulation on the active centers of Ru-based catalyst is the main reason for its deactivation, while the Fe additive can reduce the CO amount to a minimum level through WGS and FTS reaction.



美国麦克仪器公司

美国麦克仪器公司自 1962 年成立以来始终保持着颗粒分析仪器的世界领导地位，在自动样品传递、TPD/TPR 化学吸附、表面积吸附平衡、DFT 数据处理等领域拥有多达 40 多项颗粒表征方面的专利。主要产品有全自动比表面与孔隙度分析仪、化学吸附仪、高压物理 / 化学吸附仪、压汞仪、纳米粒度与 Zeta 电位分析仪、密度仪和粒度仪等。



3Flex 全功能型多用气体吸附仪

- 高通量 - 可同时进行 3 个微孔测试，超高分析能力与超多样品分析量
- 介孔、微孔分析与蒸汽吸附、化学吸附可选
- 气动高压硬密封阀，保证无泄漏气体管理
- 全新设计的自动诊断系统，可方便地对仪器各个部件实时监控
- 精确的多歧管和传感器温度控制
- 自动选择适合的压力范围显示
- 三个 RTD 用于精确温度监控和控制

研究级高性能全自动程序升温化学吸附仪 AutoChem II 2920

- 提供精确程序升温技术进行催化剂表征，如金属分散度、活性金属表面积、活性颗粒尺寸和催化材料表面活性等
- 能进行全自动脉冲化学吸附和程序升温还原（TPR）、程序升温脱附（TPD）、程序升温氧化（TPO）和程序升温反应（TPRx）以及 BET 表面积评价
- 配套专用接口，可连接质谱、气相、红外等设备
- 蒸汽发生器选件可进行惰性载气中液体蒸汽包括有机蒸汽（例如吡啶、苯）的吸附
- CryoCooler II 选件可满足低至 -120°C 的分析温度



HPVA-II 超高压容量法气体吸附仪

- 超高压，压力可达 200bar，
- 四个分析站，四个独立的脱气站，可同时测试 4 个样品
- 宽温度范围：从 -196°C 到 500°C
- 软件升级：增加自由空间管理、基线校准、压力显示等功能
- 增加传感器类型，满足不同用户的需要，仪器适用范围更广

麦克默瑞提克（上海）仪器有限公司
地址：上海市浦东新区民生路 550 号 1505-1509 室
邮编：200135 电话：021-51085884

北京办公室
地址：北京市丰台区南四环西路 188 号总部基地 12 区 28 栋 7 层
邮编：100070 电话：010-51280918

广州办公室
地址：广州市天河区中山大道 140 号华港商务大厦 1602 室
邮编：510630 电话：020-38023057

全国服务热线电话：400-630-2202
网址：www.micromeritics.com.cn, www.micromeritics.com
微博：t.sina.com.cn/micromeritics