

<http://www.chxb.cn>

ISSN 0253-9837

CN 21-1195/O6

CODEN THHPD3

催化学报


Chinese Journal of Catalysis

主编 林励吾

Editor-in-Chief LIN Liwu

2013

Vol. 34 No. 8



ISSN 0253-9837



9 770253 983139

万方数据

中国化学会催化学会会刊

Transaction of the Catalysis Society of China



中国科学院科学出版基金资助出版

催化学报

(CUIHUA XUEBAO)

CHINESE JOURNAL OF CATALYSIS

月刊 SCI收录 2013年8月 第34卷 第8期



目次

综述

1471 (英/中)

质子交换膜燃料电池Pt纳米线电催化剂研究现状
严泽宇, 李冰, 杨代军, 马建新

研究快讯

1482 (英/中)

串联双釜连续反应装置中Ru-Co-B/ZrO₂上苯选择加氢制环己烯
孙海杰, 李帅辉, 张元馨, 江厚兵, 曲良龙, 刘寿长, 刘仲毅

研究论文

1489 (英/封面文章)

单核钌催化剂化学催化和光催化水氧化反应
姜毅, 李斐, 黄芳, 张彪彪, 孙立成

1496 (英/中)

微米及纳米丝光沸石分子筛上二甲醚羰基化反应的积碳分析
薛会福, 黄秀敏, Evert Ditzel, 展恩胜, 马猛, 申文杰

1504 (英/中)

以天然凹凸棒石为原料合成Fe/Ti-ZSM-5沸石分子筛及其催化裂化性能
周晓兆, 刘艳, 孟祥举, 申宝剑, 肖丰收

1513 (英)

可磁性回收纳米级负载型杂多酸催化剂用于水中生物活性化合物的绿色合成
Ezzat Rafice, Sara Eavani, Maryam Khodayari

1519 (英/中)

巯基功能化介孔材料高效锚定钨负载型催化剂的制备及其苯酚加氢催化性能
张嘉熙, 黄高伟, 张璋, 何群华, 黄超, 杨旭, 宋慧宇, 梁振兴, 杜丽, 廖世军

1527 (英)

Ag₂S-石墨烯/TiO₂的声化学法合成及可见光催化活性
Ze-Da Meng, Lei Zhu, Kefayat Ullah, Shu Ye, Qian Sun, Won-Chun Oh

1534 (英/中)

氧化铝载体对Pt-Sn/Al₂O₃催化剂上苯胺和苯甲醇一步合成N-苯基苯胺催化反应性能的影响
殷梦瑶, 何松波, 余正坤, 吴凯凯, 王连弟, 孙承林

1543 (英)

用于苯选择加氢的Ru/Al₂O₃-ZrO₂-NiO/堇青石蜂窝整体催化剂
王铭浩, 苏宏久, 周谨, 王树东

1551 (英/中)

纳米针状氧化镓光催化降解纯水和废水中全氟辛酸
邵田, 张彭义, 李振民, 金玲

1560 (英)

α -二亚胺钴配合物催化1,3-丁二烯高活性与高顺式-1,4选择性聚合
贾翔宇, 刘恒, 胡雁鸣, 代全权, 毕吉福, 白晨曦, 张学全

1570 (英/中)

Co₂C上CO的程序升温脱附和程序升温表面反应研究
裴彦鹏, 丁云杰, 臧娟, 宋宪根, 董文达, 朱何俊, 王涛, 陈维苗

1576 (英)

多级孔ZSM-5分子筛的合成及其在甲醇脱水制二甲醚反应中的应用
杨琦, 张海涛, 孔猛, 包秀秀, 费金华, 郑小明

1583 (英)

两性芳香亚胺环钪化合物及其Langmuir-Blodgett膜的制备、表征及催化活性
赵娜, 王飞, 周梅玲, 李铁生, 刘辉, 许文俭, 吴养洁

1589 (英/中)

分子印迹聚合物负载Fe(III)催化剂的底物识别性能
孙文庆, 谭蓉, 郑卫国, 银董红

1599 (英/中)

La交换NaY分子筛中的离子定位和迁移规律
杜晓辉, 张海涛, 李雪礼, 谭争国, 刘宏海, 高雄厚

1608 (英/中)

界面活化的溶胶凝胶包埋Candida rugosa脂肪酶催化合成维生素E琥珀酸酯
胡焱, 蒋相军, 吴素文, 江凌, 黄和

1617 (英)

硫掺杂橄榄状BiVO₄上可见光降解亚甲基蓝和甲醛水溶液性能
赵振璇, 戴洪兴, 邓积光, 刘雨溪, 区泽棠


1627 (英)

仿生制备有机-无机复合微囊固定化葡萄糖氧化酶
辛茜, 姜艳军, 高静, 周丽亚, 马丽, 贺莹, 贾霏


相关信息

1634 作者索引


英文全文电子版(国际版)由Elsevier出版社在ScienceDirect上出版
<http://www.sciencedirect.com/science/journal/18722067>
<http://www.elsevier.com/locate/chnjc>
<http://www.chxb.cn>



available at www.sciencedirect.com



journal homepage: www.elsevier.com/locate/chnjc



Chinese Journal of Catalysis


Graphical Contents

Review

Chin. J. Catal., 2013, 34: 1471–1481 doi: 10.1016/S1872-2067(12)60629-9

Pt nanowire electrocatalysts for proton exchange membrane fuel cells

Zeyu Yan, Bing Li*, Daijun Yang, Jianxin Ma
Tongji University



This review presents the research progress of different preparation methods of Pt nanowire catalyst and its electrochemical performance, durability, and CO tolerance applying to proton exchange membrane fuel cells.


Communication

Chin. J. Catal., 2013, 34: 1482–1488 doi: 10.1016/S1872-2067(12)60637-8

Selective hydrogenation of benzene to cyclohexene in continuous reaction device with two reaction reactors in serie over Ru-Co-B/ZrO₂ catalysts

Haijie Sun, Shuaihui Li, Yuanxin Zhang, Houbing Jiang, Lianglong Qu,
Shouchang Liu, Zhongyi Liu*
Zhengzhou University;
Zhengzhou Normal University;
Beijing Energy Engineering Technologies Co., Ltd

The selectivity for cyclohexene and the cyclohexene yield were stabilized at around 73% and 30%, respectively, in 419 h over a Ru-Co-B/ZrO₂ catalyst in a continuous device with two reactors in series.




Articles

Chin. J. Catal., 2013, 34: 1489–1495 doi: 10.1016/S1872-2067(12)60600-7

Chemical and photocatalytic water oxidation by mononuclear Ru catalysts

Yi Jiang, Fei Li*, Fang Huang, Biaobiao Zhang, Licheng Sun*
Dalian University of Technology, China;
KTH Royal Institute of Technology, Sweden


Mononuclear ruthenium complexes based on the bipyridine-dicarboxylate (bda) ligand were prepared and showed high catalytic efficiencies for chemical and photochemical water oxidation, a key challenge for solar energy conversion into fuels. The more electron-withdrawing substituents on the axial ligands of the catalysts lead to higher activities towards water oxidation.



Chin. J. Catal., 2013, 34: 1496–1503 doi: 10.1016/S1872-2067(12)60607-X

Coking on micrometer- and nanometer-sized mordenite during dimethyl ether carbonylation to methyl acetate

Huifu Xue, Xiumin Huang, Evert Ditzel, Ensheng Zhan, Meng Ma, Wenjie Shen*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences, China; BP Chemicals Limited, UK




Nanometer-sized H-mordenite (HMOR) showed a much higher reaction rate and stability than micrometer-sized HMOR, resulting from the improvement of molecule transportation efficiency by reducing the crystalline size.

Chin. J. Catal., 2013, 34: 1504–1512 doi: 10.1016/S1872-2067(12)60638-X

Synthesis and catalytic cracking performance of Fe/Ti-ZSM-5 zeolite from attapulgite mineral

Xiaozhao Zhou, Yan Liu, Xiangju Meng, Baojian Shen, Feng-Shou Xiao*
Zhejiang University; China University of Petroleum




Fe/Ti-ZSM-5 zeolite was synthesized from treated attapulgite (ATP) mineral. Compared with conventional ZSM-5 zeolite, Fe/Ti-ZSM-5 exhibits relatively high yields of light olefins in the catalytic cracking of Canadian light gas oil.

Chin. J. Catal., 2013, 34: 1513–1518 doi: 10.1016/S1872-2067(12)60645-7

Magnetically recoverable, nanoscale-supported heteropoly acid catalyst for green synthesis of biologically active compounds in water

Ezzat Rafiee*, Sara Eavani, Maryam Khodayari
Razi University, Iran




12-Tungstophosphoric acid (PW) catalysts supported on aerosil silica and silica-coated γ - Fe_2O_3 nanoparticles ($\text{Fe}_2\text{O}_3@SiO_2$ -PW) were prepared and characterized. These catalysts were used for the aqueous syntheses of 1,8-dioxo-9,10-diaryldecahydroacridines and xanthene derivatives, which are biologically interesting compounds.

Chin. J. Catal., 2013, 34: 1519–1526 doi: 10.1016/S1872-2067(12)60603-2

Immobilization of highly active Pd nano-catalysts on functionalized mesoporous silica supports using mercapto groups as anchoring sites and their catalytic performance for phenol hydrogenation

Jiayi Zhang, Gaowei Huang, Cheng Zhang, Qunhua He, Chao Huang,
Xu Yang, Huiyu Song, Zhenxing Liang, Li Du*, Shijun Liao*
South China University of Technology;
Guangdong Environmental Monitoring Center;
Guangzhou Institute of Energy Conversion, Chinese Academy of Sciences


Highly dispersed Pd nanoparticle catalyst on -SH-functionalized mesoporous silica (Pd-SH-MCM-41) was prepared by anchoring interaction between -SH groups and Pd cations. This catalyst showed very high catalytic activity (>99%) for the hydrogenation of phenol.



Chin. J. Catal., 2013, 34: 1527–1533 doi: 10.1016/S1872-2067(12)60611-1

Enhanced visible light photocatalytic activity of Ag_2S -graphene/ TiO_2 nanocomposites made by sonochemical synthesis

Ze-Da Meng, Lei Zhu, Kefayat Ullah, Shu Ye, Qian Sun, Won-Chun Oh*
Hanseu University, Korea




This paper presents the research of synthesized Ag_2S -graphene/ TiO_2 by the sonochemical method. The generation of reactive oxygen species was detected through the oxidation reaction from 1,5-diphenyl carbazide to 1,5-diphenyl carbazone.

Chin. J. Catal., 2013, 34: 1534–1542 doi: 10.1016/S1872-2067(12)60608-1

Effect of alumina support on catalytic performance of Pt-Sn/Al₂O₃ catalysts in one-step synthesis of *N*-phenylbenzylamine from aniline and benzyl alcohol

Mengyao Yin, Songbo He*, Zhengkun Yu, Kaikai Wu, Liandi Wang, Chenglin Sun*

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences




Pt-Sn/Al₂O₃ catalysts with highly dispersed Pt particles, weak acid sites, acid distributions, and large pore volumes and pore size distributions have excellent catalytic performance in the synthesis of secondary amines.

Chin. J. Catal., 2013, 34: 1543–1550 doi: 10.1016/S1872-2067(12)60609-3

Ru/Al₂O₃-ZrO₂-NiO/cordierite monolithic catalysts for selective hydrogenation of benzene

Minghao Wang, Hongjiu Su, Jin Zhou, Shudong Wang*

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences




Novel egg-shell-like Ru/Al₂O₃-ZrO₂-NiO/cordierite monolithic catalysts with tailored pore structure are fabricated. Mesopores and low pore volume are needed to obtain a reasonable yield of cyclohexene using the monolithic catalysts with a low concentration of ZnSO₄ solution.

Chin. J. Catal., 2013, 34: 1551–1559 doi: 10.1016/S1872-2067(12)60612-3

Photocatalytic decomposition of perfluorooctanoic acid in pure water and wastewater by needle-like nanostructured gallium oxide

Tian Shao, Pengyi Zhang*, Zhenmin Li, Ling Jin

Tsinghua University




In combination with 185 nm vacuum UV irradiation, needle-like Ga₂O₃ showed high efficiency for the removal of trace perfluorooctanoic acid (PFOA) in wastewater where the decomposition of PFOA by other catalysts is usually inhibited by coexisting natural organic matters.

Chin. J. Catal., 2013, 34: 1560–1569 doi: 10.1016/S1872-2067(12)60625-1

Highly active and *cis*-1,4 selective polymerization of 1,3-butadiene catalyzed by cobalt(II) complexes bearing α -diimine ligands

Xiangyu Jia, Heng Liu, Yanming Hu*, Quanquan Dai, Jifu Bi, Chenxi Bai, Xuequan Zhang*
 Changchun Institute of Applied Chemistry, Chinese Academy of Sciences; University of Chinese Academy of Sciences;
 Dalian University of Technology




A series of cobalt(II) complexes ligated by α -diimine have been synthesized and characterized. The catalyst system [α -diimine]CoCl₂/EASC shows high catalytic activity and *cis*-1,4 selectivity (up to 98%) for 1,3-butadiene polymerization.

Chin. J. Catal., 2013, 34: 1570–1575 doi: 10.1016/S1872-2067(12)60615-9

Temperature-programmed desorption and surface reaction studies of CO on Co₂C

Yanpeng Pei, Yunjie Ding*, Juan Zang, Xiangen Song, Wenda Dong, Hejun Zhu, Tao Wang, Weimiao Chen
 Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences




Co₂C was prepared by carburizing Co with CO. The prepared Co₂C samples were composed of a bulk Co₂C phase and an outer CoO passivation layer. Co₂C could adsorb CO, which was hydrogenated into alcohol.

Chin. J. Catal., 2013, 34: 1576–1582 doi: 10.1016/S1872-2067(12)60621-4

Hierarchical mesoporous ZSM-5 for the dehydration of methanol to dimethyl ether

Qi Yang, Haitao Zhang, Meng Kong, Xiuxiu Bao, Jinhua Fei*, Xiaoming Zheng
 Zhejiang University; Petrochina Lanzhou Chemical Research Centre




The synthesized hierarchical mesoporous HZSM-5 zeolites exhibited better catalytic activity than conventional HZSM-5, indicating that hierarchical mesopores play an important role in the methanol to dimethyl ether process.

Chin. J. Catal., 2013, 34: 1583–1588 doi: 10.1016/S1872-2067(12)60613-5

Preparation, characterization and catalytic activity of amphiphilic cyclopalladated aryl imines and their Langmuir-Blodgett films

Na Zhao, Fei Wang, Meiling Zhou, Tiesheng Li*, Hui Liu, Wenjian Xu, Yangjie Wu*
Zhengzhou University




The amphiphilic cyclopalladated aryl imine catalysts were successfully transferred on the solid slides and can be used for the Suzuki reactions.

Chin. J. Catal., 2013, 34: 1589–1598 doi: 10.1016/S1872-2067(12)60624-X

Molecularly imprinted polymer containing Fe(III) catalysts for specific substrate recognition

Wenqing Sun, Rong Tan*, Weiguo Zheng, Donghong Yin*
Hunan Normal University;
Tobacco Hunan Industrial Corporation


A series of molecularly imprinted polymers (MIPs) containing the equal amount of iron(III) were found to be the specific substrate recognition catalysts in the oxidation of substituted benzyl alcohol in water.



Chin. J. Catal., 2013, 34: 1599–1607 doi: 10.1016/S1872-2067(12)60622-6

Cation location and migration in lanthanum-exchanged NaY zeolite

Xiaohui Du, Haitao Zhang, Xueli Li, Zhengguo Tan, Honghai Liu, Xionghou Gao*
Northwest Normal University; Lanzhou Petrochemical Research Center




Lanthanum cations are initially distributed in supercages and then migrate to SI' located inside sodalite cages during heating and dehydration.

Chin. J. Catal., 2013, 34: 1608–1616 doi: 10.1016/S1872-2067(12)60628-7

Synthesis of vitamin E succinate by interfacial activated *Candida rugosa* lipase encapsulated in sol-gel materials

Yi Hu, Xiangjun Jiang, Suwen Wu, Ling Jiang, He Huang*
Nanjing University of Technology




Interfacial activated *Candida rugosa* lipase encapsulated in sol-gel materials was prepared and used to synthesize vitamin E succinate.

Chin. J. Catal., 2013, 34: 1617–1626 doi: 10.1016/S1872-2067(12)60632-9

Effect of sulfur doping on the photocatalytic performance of BiVO_4 under visible light illumination

Zhenxuan Zhao, Hongxing Dai*, Jiguang Deng, Yuxi Liu, Chak Tong Au*
Beijing University of Technology; Hong Kong Baptist University




Porous $\text{BiVO}_{4-x}\text{S}_x$ and $\text{BiVO}_{4-x}\text{S}_x$ are fabricated using dodecylamine-assisted alcohol-hydrothermal strategy. The higher O_{ads} concentration and lower bandgap energy account for excellent photocatalytic performance of $\text{BiVO}_{4-x}\text{S}_{0.08}$ for methylene blue and formaldehyde degradation.

Chin. J. Catal., 2013, 34: 1627–1633 doi: 10.1016/S1872-2067(12)60635-4

Biomimetic preparation of organic-inorganic composite microcapsules for glucose oxidase immobilization

Qian Xin, Yanjun Jiang*, Jing Gao*, Liya Zhou, Li Ma, Ying He, Fei Jia
Hebei University of Technology; Institute of Process Engineering, Chinese Academy of Sciences



Glucose oxidase (GOD) was immobilized in organic-inorganic composite microcapsules using a combination of layer-by-layer assembly and biomimetic mineralization. The encapsulated GOD exhibited distinct advantages in terms of thermal, pH, and operational stabilities.