



Chinese Journal of Catalysis


www.cjcatal.org

Volume 37 | Number 1 | January 2016

新型多孔催化材料专栏

Special Column on New Porous Catalytic Materials

Guest Editors: Ying Wan (万颖), Serge Kaliaguine,
Dongyuan Zhao (赵东元)




Science Press
万方数据




Editors-in-Chief Can Li (李灿) Tao Zhang (张涛)
Transaction of The Catalysis Society of China



available at www.sciencedirect.com



journal homepage: www.elsevier.com/locate/chnjc



Chinese Journal of Catalysis

Graphical Contents

Special Column on New Porous Catalytic Materials

Chin. J. Catal., 2016, 37: 1–2 doi: 10.1016/S1872-2067(15)61027-0

Preface to Special Column on New Porous Catalytic Materials

Ying Wan, Serge Kaliaguine, Dongyuan Zhao (Guest Editors)


Shanghai Normal University, China; Université Laval, Canada; Fudan University, China

*Chin. J. Catal.*, 2016, 37: 3–15 doi: 10.1016/S1872-2067(15)60901-9 [Review]

Ethylbenzene to styrene over ZrO₂ based mixed metal oxide catalysts with CO₂ as a soft oxidant

Nanzhe Jiang, Abhishek Burri, Sang-Eon Park*

Yanbian University, China; Inha University, Korea




Acid-base bi-functionality activation of alkyl aromatics and CO₂ is simultaneously performed on the same zirconia-based mixed metal oxide catalyst to yield high activity and selectivity.

Chin. J. Catal., 2016, 37: 16–26 doi: 10.1016/S1872-2067(15)60965-2 [Review]

Advances in development and industrial applications of ethylbenzene processes

Weimin Yang*, Zhendong Wang, Hongmin Sun, Bin Zhang
Sinopec Shanghai Research Institute of Petrochemical Technology




Advances in catalysts and processes for benzene ethylation with pure/dilute ethylene or ethanol to produce ethylbenzene are summarized. Some promising techniques and applications are discussed.

Chin. J. Catal., 2016, 37: 27–31 doi: 10.1016/S1872-2067(15)61026-9 [Communication]

Synthesis of three-dimensional ordered mesoporous MnO₂ and its catalytic performance for formaldehyde oxidation

Bingyang Bai, Qi Qiao*, Junhua Li*, Jiming Hao
Chinese Research Academy of Environmental Sciences;
Tsinghua University


Three-dimensional ordered mesoporous MnO₂ was synthesized using KIT-6 molecule sieves as a hard template. The material showed high catalytic activity in formaldehyde oxidation and has good development potential.



Chin. J. Catal., 2016, 37: 32–42 doi: 10.1016/S1872-2067(15)60909-3 [Article]

Pore structure effects on the kinetics of methanol oxidation over nanocast mesoporous perovskites

Mahesh M. Nair, Freddy Kleitz, Serge Kaliaguine*
Université Laval, Canada




Changing SBA-15 template aging temperature which affects the pore structure of the mesoporous perovskite does not change activation energy. Only preexponential factor is changed proportionally to specific surface area.

Chin. J. Catal., 2016, 37: 43–53 doi: 10.1016/S1872-2067(15)60878-6 [Article]

Ordered mesoporous carbon supported bifunctional PtM (M = Ru, Fe, Mo) electrocatalysts for a fuel cell anode

Chin-Te Hung, Zih-Hao Liou, Pitchaimani Veerakumar, Pei-Hao Wu, Tuan-Chi Liu*, Shang-Bin Liu*

Institute of Atomic and Molecular Sciences, Academia Sinica, Taiwan, China; National Taiwan University of Science and Technology, Taiwan, China; National Taiwan Normal University, Taiwan, China




The deposition onto an ordered mesoporous carbon (OMC) support of well dispersed PtM (M = Ru, Fe, Mo) alloy nanoparticles (NPs) was directly replicated. These PtM-OMC catalysts use a cost-effective second metal such as Fe and possess uniform (2–3 nm) and highly dispersed alloy NPs catalysts that exhibit superior electrocatalytic performance and excellent tolerance to CO poisoning during the methanol oxidation reaction.

Chin. J. Catal., 2016, 37: 54–60 doi: 10.1016/S1872-2067(15)60952-4 [Article]

A Pd-metalated porous organic polymer as a highly efficient heterogeneous catalyst for C-C couplings

Zhifeng Dai, Fang Chen, Qi Sun, Yanyan Ji, Liang Wang, Xiangju Meng*, Feng-Shou Xiao

Zhejiang University



A Pd-metalated porous polymer bearing phenanthroline moieties (Pd/POP-Phen) exhibits high activities, excellent selectivities and superior recyclabilities in Suzuki, Sonogashira and Heck couplings.

Chin. J. Catal., 2016, 37: 61–72 doi: 10.1016/S1872-2067(15)60917-2 [Article]

Size-control growth of thermally stable Au nanoparticles encapsulated within ordered mesoporous carbon framework

Shuai Wang, Jie Wang, Xiaojuan Zhu, Jianqiang Wang,

Osamu Terasaki, Ying Wan*


Shanghai Normal University, China;

Shanghai Institute of Applied Physics,

Chinese Academy of Sciences, China;

Stockholm University, Sweden


Thermally stable gold nanoparticles with sizes of 3–18 nm and free of protecting agent have been loaded onto ordered mesoporous carbon via a coordination-assisted self-assembly approach.



Chin. J. Catal., 2016, 37: 73–82 doi: 10.1016/S1872-2067(15)60951-2 [Article]

Rh₂O₃/mesoporous MO_x-Al₂O₃ (M = Mn, Fe, Co, Ni, Cu, Ba) catalysts: Synthesis, characterization, and catalytic applications

Huan Liu, Yi Lin, Zhen Ma*
Fudan University




Mesoporous MO_x-Al₂O₃ (M = Mn, Fe, Co, Ni, Cu, Ba) materials were prepared by a one-pot evaporation-induced self-assembly method. Rh₂O₃ nanoparticles were loaded onto these mesoporous supports via wet impregnation.

Chin. J. Catal., 2016, 37: 83–90 doi: 10.1016/S1872-2067(15)60913-5 [Article]

Synthesis of Pd nanoparticles supported on CeO₂ nanotubes for CO oxidation at low temperatures

Jingmi Wu, Liang Zeng, Dangguo Cheng*, Fengqiu Chen, Xiaoli Zhan, Jinlong Gong*
Zhejiang University; Tianjin University




The preparation of CeO₂ nanotube-supported Pd nanoparticles and their catalytic performance toward CO oxidation were described. Abundant Ce³⁺ and Pd–O–Ce interactions on the nanotube facilitated the reaction.

Chin. J. Catal., 2016, 37: 91–97 doi: 10.1016/S1872-2067(15)60940-8 [Article]

Ni@Pd core-shell nanoparticles supported on a metal-organic framework as highly efficient catalysts for nitroarenes reduction

Siping Jian, Yingwei Li*
South China University of Technology



Bimetal core-shell Ni@Pd nanoparticles supported on MIL-101 were prepared and exhibit high catalytic activity and selectivity during the reduction of nitroarenes.


Highlight

Chin. J. Catal., 2016, 37: 98–101 doi: 10.1016/S1872-2067(15)61008-7

Recent advances of remote selective C–H activation: Ligand and template design

Guoqiang Yang, Nicholas Butt, Wanbin Zhang*
Shanghai Jiao Tong University

remote site selective C–H activation: template and ligand design




The *para*-selective C–H activation of toluene-type substrates has been achieved using a strained biphenyl-type template, and a highly selective *meta*-C–H borylation of aromatic amides has been reported by taking advantage of secondary ligand interactions.

Review

Chin. J. Catal., 2016, 37: 102–122 doi: 10.1016/S1872-2067(15)61007-5

Progress in research on catalysts for catalytic oxidation of formaldehyde

Bingyang Bai*, Qi Qiao, Junhua Li*, Jiming Hao
Chinese Research Academy of Environmental Sciences; Tsinghua University



This paper reviews progress in research on precious-metal and transition-metal oxide catalyst systems for HCHO oxidation. The oxidation properties, factors influencing the catalytic activity, and reaction mechanisms are discussed, and future development directions and research hotspots are considered.


Articles

Chin. J. Catal., 2016, 37: 123–134 doi: 10.1016/S1872-2067(15)61012-9

Flower-like 3D CuO microsphere acting as photocatalytic water oxidation catalyst

Xiaoqiang Du, Jingwei Huang, Yingying Feng, Yong Ding*
Lanzhou University;
Lanzhou Institute of Chemical Physics,
Chinese Academy of Sciences


A flower-like 3D CuO microspheres based on earth abundant elements has been successfully synthesized. It is the first time that a copper oxide shows efficient photocatalytic water oxidation performance with good stability.



Chin. J. Catal., 2016, 37: 135–145 doi: 10.1016/S1872-2067(15)61010-5

Oxidation of 4-chloro-3-methylphenol using zeolite Y-encapsulated iron(III)-, nickel(II)-, and copper(II)-*N,N'*-disalicylidene-1,2-phenylenediamine complexes

Solomon Legese Hailu, Balachandran Unni Nair *, Mesfin Redi-Abshiro, Isabel Diaz, Rathinam Aravindhan, Merid Tessema
CSIR-Central Leather Research Institute, India; Addis Ababa University, Ethiopia; Instituto de Catálisis y Petroleoquímica, Spain




Iron(III), nickel(II) and copper(II)-*N,N'*-disalicylidene-1,2-phenylenediamine complex is successfully encapsulated through ship-in-a-bottle method into zeolite Y supercage. Zeolite encapsulated metal complexes show good catalytic efficiency towards degradation of 4-chloro-3-methylphenol.

Chin. J. Catal., 2016, 37: 146–152 doi: 10.1016/S1872-2067(15)61005-1

Ultrasound assisted multicomponent reactions: A green method for the synthesis of *N*-substituted 1,8-dioxo-decahydroacridines using β -cyclodextrin as a supramolecular reusable catalyst in water

Asha V. Chate, Umesh B. Rathod, Jagdish S. Kshirsagar, Pradip A. Gaikwad, Kishor D. Mane, Pravin S. Mahajan, Mukesh D. Nikam, Charansingh H. Gill *
Dr. Babasaheb Ambedkar Marathwada University, India




A mild and environmentally benign protocol was developed using β -cyclodextrins as a catalyst to significantly improve the yield of *N*-substituted 1,8-dioxo-decahydroacridine derivatives. The β -cyclodextrin employed in the multi-component reaction satisfies the requirements for green chemistry.

Chin. J. Catal., 2016, 37: 153–158 doi: 10.1016/S1872-2067(15)61001-4

Ruthenium nanoparticles supported on mesoporous MCM-41 as an efficient and reusable catalyst for selective oxidation of arenes under ultrasound irradiation

Alireza Khorshidi *
University of Guilan, Iran




MCM-41 crystallites loaded with Ru nanoparticles were used as heterogeneous catalysts in selective oxidation of arenes under sonic waves. The catalyst active site was found to be RuO_2 species formed during the course of reaction.

Chin. J. Catal., 2016, 37: 159–168 doi: 10.1016/S1872-2067(15)60990-1

Preparation of a novel supported electrode comprising a nickel (II) hydroxide-modified carbon paste electrode ($\text{Ni}(\text{OH})_2\text{-X/CPE}$) for the electrocatalytic oxidation of formaldehyde

Safura Kaviani, Seyed Naser Azizi *, Shahram Ghasemi
University of Mazandaran, Iran

A novel $\text{Ni}(\text{OH})_2\text{-X/CPE}$ was first prepared and used for the electrocatalytic oxidation of formaldehyde. This electrode exhibited good electrocatalytic activity with regards to formaldehyde oxidation owing to its nanoporous structure and the large surface area of zeolite X.




Chin. J. Catal., 2016, 37: 169–176 doi: 10.1016/S1872-2067(15)60982-2

Comparison of growth mechanisms of undoped and nitrogen-doped carbon nanofibers on nickel-containing catalysts

Vladimir V. Chesnokov, Olga Yu. Podyacheva *, Alexander N. Shmakov,
Lidiya S. Kibis, Andrei I. Boronin, Zinfer R. Ismagilov
Boreskov Institute of Catalysis, Russia;
Novosibirsk State University, Russia;
Institute of Coal Chemistry and Materials Science, Russia

A similarity of the growth mechanisms of carbon nanofibers and nitrogen-doped carbon nanofibers on Ni-containing catalysts was demonstrated by the use of XPS and XRD in situ techniques.




Chin. J. Catal., 2016, 37: 177–183 doi: 10.1016/S1872-2067(15)60976-7

Effects of alkaline additives on the formation of lactic acid in sorbitol hydrogenolysis over Ni/C catalyst

Junjie Zhang, Fang Lu *, Weiqiang Yu, Rui Lu, Jie Xu *

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences




La(OH)₃ has been identified as an efficient additive for sorbitol hydrogenolysis over Ni/C with very little lactic acid formation. The pathways for 1,2-propylene glycol and lactic acid formation were competitive, with the hydrogenation to 1,2-propylene glycol dominating for La(OH)₃.

Chin. J. Catal., 2016, 37: 184–192 doi: 10.1016/S1872-2067(15)60983-4

Solvent-free selective oxidation of cyclohexane with molecular oxygen over manganese oxides: Effect of the calcination temperature

Mingzhou Wu, Wangcheng Zhan *, Yun Guo, Yunsong Wang, Yanglong Guo, Xueqing Gong, Li Wang, Guanzhong Lu *

East China University of Science and Technology



MnO_x exhibited high catalytic activity and excellent stability during cyclohexane oxidation with molecular oxygen. The MnO_x phase state, as determined by the calcination temperature, had a crucial effect on its catalytic activity.


Chin. J. Catal., 2016, 37: 193–198 doi: 10.1016/S1872-2067(15)60969-X

Theoretical studies of CO oxidation with lattice oxygen on Co₃O₄ surfaces

Yang-Gang Wang, Xiao-Feng Yang, Jun Li *

Tsinghua University;
Dalian Institute of Chemical Physics,
Chinese Academy of Sciences

Theoretical studies reveal that the reactivity of CO oxidation on Co₃O₄ nanocrystals exhibits significant crystal plane effects and structure sensitivity, and has a positive correlation with the surface reducibility.





中国科学院科学出版基金资助出版



Chinese Journal of
Catalysis

月刊 SCI 收录 2016 年 1 月 第 37 卷 第 1 期



目 次

新型多孔催化材料专栏

1 (专栏前言)

新型多孔催化材料专栏前言

万颖, Serge Kaliaguine, 赵东元 (客座主编)

3 (综述)

ZrO₂ 基复合金属氧化物催化剂上 CO₂ 温和氧化乙苯制苯乙烯

Nanzhe Jiang, Abhishek Burri, Sang-Eon Park

16 (综述)

乙苯工艺技术开发及工业应用进展

杨为民, 王振东, 孙洪敏, 张斌

27 (快讯)

三维有序介孔二氧化锰制备及其甲醛催化氧化性能

拜冰阳, 乔琦, 李俊华, 郝吉明

32 (论文)

孔结构对纳米浇注的中孔钙钛矿上甲醇氧化反应动力学的影响

Mahesh M. Nair, Freddy Kleitz, Serge Kaliaguine

43 (论文)

用于燃料电池阳极的有序介孔碳负载的双功能 Pt-M (M = Ru, Fe, Mo) 电催化剂

洪锦德, 刘子豪, 维拉库玛, 吴培豪, 刘端祺, 刘尚斌

54 (论文)

多孔有机聚合物负载钨作为高效 C-C 偶联反应多相催化剂

戴志锋, 陈芳, 孙琦, 纪妍妍, 王亮, 孟祥举, 肖丰收

61 (论文)

尺寸可控、高热稳定性有序介孔碳 Au 催化剂的合成

王帅, 王杰, 朱小娟, 王建强, Osamu Terasaki, 万颖

73 (论文)

Rh₂O₃/介孔 MO_x-Al₂O₃ (M = Mn, Fe, Co, Ni, Cu, Ba) 催化剂: 合成、表征和催化应用

刘欢, 林毅, 马臻

83 (论文)

CeO₂ 纳米管负载 Pd 纳米颗粒催化 CO 低温氧化

吴静谧, 曾亮, 程党国, 陈丰秋, 詹晓力, 巩金龙

91 (论文)

MIL-101 负载 Ni@Pd 核壳纳米粒子催化芳香硝基类化合物加氢

简思平, 李映伟

亮 点

98

远程选择性碳氢活化的近期进展: 模板与配体设计

杨国强, 白尼克, 张万斌

综 述

102

甲醛催化氧化催化剂的研究进展

拜冰阳, 乔琦, 李俊华, 郝吉明

论 文

123

花状的三维 CuO 微球作为光催化水氧化催化剂

杜晓强, 黄静伟, 丰营营, 丁勇

135

封装 Fe(III), Ni(II) 和 Cu(II) 的 *N,N'*-双邻羟苯亚甲基-1,2-苯二胺络合物的 Y 分子筛催化 4-氯-3-甲基苯酚氧化反应

Solomon Legese Hailu, Balachandran Unni Nair, Mesfin Redi-Abshiro, Isabel Diaz, Rathinam Aravindhan, Merid Tessema

146

超声辅助多组分反应: 水介质中以 β-环糊精为可重复使用的超分子催化剂绿色合成氮取代的 1,8-二氧-十氢吡啶

Asha V. Chate, Umesh B. Rathod, Jagdish S. Kshirsagar, Pradip A. Gaikwad, Kishor D. Mane, Pravin S. Mahajan, Mukesh D. Nikam, Charansingh H. Gill

153

中孔 MCM-41 负载 Ru 纳米粒子催化剂用于超声辐射下芳烃选择氧化反应

Alireza Khorshidi

159

用于电催化氧化甲醛的新型 Ni(OH)₂-X/CPE 负载电极的制备

Safura Kaviani, Seyed Naser Azizi, Shahram Ghasemi

169

Ni 基催化剂上未掺杂和氮掺杂的碳纳米纤维生长机理的比较

Vladimir V. Chesnokov, Olga Yu. Podyacheva, Alexander N. Shmakov, Lidiya S. Kibis, Andrei I. Boronin, Zinifer R. Ismagilov

177

Ni/C 催化剂上山梨醇氢解反应中碱对乳酸形成的影响
张俊杰, 路芳, 于维强, 卢锐, 徐杰

184

氧化锰催化剂在环己烷无溶剂选择性氧化反应中的活性:
焙烧温度的影响

吴明周, 詹望成, 郭耘, 王筠松, 郭杨龙, 龚学庆, 王丽,
卢冠忠

193

Co₃O₄ 纳米晶表面晶格氧催化氧化 CO 的理论研究
王阳刚, 杨小峰, 李隽

相关信息

199 《催化学报》第五届编辑委员会

202 Guide for Authors

209 《催化学报》作者指南

英文全文电子版(国际版)由Elsevier出版社在ScienceDirect上出版

<http://www.sciencedirect.com/science/journal/18722067>

<http://www.elsevier.com/locate/chnjc>

www.cjcatal.org

在线投审稿网址


<https://mc03.manuscriptcentral.com/cjcatal>

美国麦克仪器公司-材料表征权威




美国麦克仪器成立于1962年，是材料特性实验室分析仪器和服务的领导者。公司致力于生产分析粉末/固体材料物理化学性质的全自动化仪器，能够进行比表面积、孔容、孔径、孔径分布、密度、催化剂性质表征、催化剂活性测试以及粒度粒形分析，可广泛用于基础研究、产品开发及质量控制等各个阶段。

美国麦克仪器公司早在1979年就进入中国市场，是中美建交后最早进入中国市场的分析仪器。在为中国用户服务30多年后，于2011年3月在上海成立了麦克默瑞提克（上海）仪器有限公司，专业为中国市场提供美国麦克仪器公司的产品和服务。2014年8月，公司在上海成立大型分析服务中心，提供全面的分析测试服务。



ASAP®
2460系列

多站扩展式全自动比表面与孔隙度分析仪



3Flex系列

三站多功能性多用气体吸附仪



Microactivity
Effi 系列

催化剂评价整体方案



AutoChem
系列

高性能全自动化学吸附仪



扫描关注官方微信

万方数据

麦克默瑞提克（上海）仪器有限公司

地址：上海市民生路600号船研大厦1505-1509室 电话：021-51085884

邮编：200135

全国服务热线电话：400-630-2202

ISSN 0253-9837

