



Chinese Journal of Catalysis


www.cjatal.com

Volume 43 | Number 7 | July 2022

二氧化碳催化转化专栏

Special Column on Catalytic Conversion of CO₂

Guest Editors: Yanqiang Huang (黄延强), Da-Gang Yu (余达刚), Guoxiong Wang (汪国雄)



催化学报
CHINESE JOURNAL OF CATALYSIS

July 2022

Vol. 43 No. 7

Pages 1545-1962

科学出版社



CHINESE
CHEMICAL
SOCIETY

Editors-in-Chief Can Li Tao Zhang
Transaction of The Catalysis Society of China

目次

二氧化碳催化转化专栏

1545 (专栏前言)

二氧化碳催化转化专栏

黄延强, 余达刚, 汪国雄 (客座主编)

1547 (综述)

单原子基催化剂用于电化学CO₂还原

孙倩, 贾忱, 赵勇, 赵川

1598 (综述)

多组分反应策略将二氧化碳固定为氨基甲酸酯的研究进展

汪露, 戚朝荣, 熊文芳, 江焕峰

1618 (综述)

二氧化碳电还原催化剂调控策略与结构效应

刘聪, 梅轩豪, 韩策, 宫雪, 宋平, 徐维林

1634 (综述)

二氧化碳电还原为甲烷的研究进展

郑汉, 杨正午, 孔祥栋, 耿志刚, 曾杰

1642 (论文)

基于碳二亚胺调控的甲酸原位产生一氧化碳策略的炔烃氢羧化反应: 二氧化碳的间接利用

夏书梅, 杨志文, 陈凯宏, 王宁, 何良年

1652 (快讯)

光催化苄基邻卤芳醚与CO₂的连续去芳构化/羧化反应合成螺环羧酸

易雅平, 席婵娟

1657 (论文)

梯型2D/2D Bi₂MoO₆/BiOI范德华异质结的制备及CO₂光还原性能

王中辽, 程蓓, 张留洋, 余家国, 李佑稷, S. Wageh, Ahmed A. Al-Ghamdi

1667 (快讯)

可见光催化二氧化碳与苄位卤代物的羧基化: 温和条件且无过渡金属

敬科, 魏明恺, 颜思顺, 廖黎丽, 牛亚楠, 罗书平, 于博, 余达刚

1674 (快讯)

调节In@InO_x核壳纳米颗粒中的表面In-O提升电催化还原CO₂为甲酸盐性能

杨艳, 傅佳驹, 唐堂, 牛帅, 张礼兵, 张佳楠, 胡劲松

1680 (快讯)

通过金属间化合物PdBi纳米片调控Pd基催化剂电催化CO₂还原为甲酸盐

谢林峰, 刘轩, 黄凡洋, 梁嘉顺, 刘健云, 王谭源, 杨利明, 曹瑞国, 李菁

1687 (论文)

金属有机框架-离子液体混合催化剂用于电化学还原二氧化碳生成甲烷

Ernest Pahuyo Delmo, 王忆安, 王菁, 朱尚乾, 李铁怀, 秦雪苹, 田一博, 赵青蓝, Juhee Jang, 王一诺, 谷猛, 张莉莉, 邵敏华

1697 (论文)

氮掺杂碳催化剂在高效电化学CO₂还原中的协同效应

刘储浩, 吴悦, 方锦杰, 余珂, 李慧, 贺文君, 张永臻, 柳守杰, 陈郑, 董京, 陈晨

1703 (论文)

类空气浓度的二氧化碳的高效电还原

陈杨岫, 阚森, 燕帅, 张俊波, 刘坤豪, 严雅琴, 关安翔, 吕希蒙, 钱林平, 郑耿锋

1710 (论文)

镍高度分散在铁基钙钛矿用于高温固体氧化物电解池阴极的CO₂电催化还原研究

周莹杰, 刘天夫, 宋月锋, 吕厚甫, 刘清雪, 塔娜, 张小敏, 汪国雄

综 述

1719

环保型碳纳米材料在光催化制氢方面的应用

熊胜, 汤榕葭, 龚道新, 邓焱成, 郑蒋夫, 李玲, 周展鹏, 杨丽华, 苏龙

1749

无细胞多酶系统的模块化构建

张寅晨, 聂宁, 张一飞

1761

光电化学氮还原: 一种可持续的氨合成方法

王立群, 闫啸, 司文平, 刘道兰, 侯兴刚, 李德军, 侯峰, 窦世学, 梁骥

1774

光催化水分解中地球储量丰富助催化剂的光沉积方法、功能与机理

赵辉, 茅沁怡, 蹇亮, 董玉明, 朱永法

快 讯

1805

有机-无机杂化钙钛矿基光催化剂用于可见光催化芳香族溴化反应并联合产氢气
张艳飞, 王洪, 刘葵, 李灿

1812

钼(0)/铜(I)协同催化脱羧不对称4+2环加成反应合成四环香豆素
王凯, 林祥丰, 李倩, 刘葵, 李灿

论 文

1818

双助催化剂负载CdS的能量捕集和电荷分离增强产H₂
张美玉, 秦朝朝, 孙万军, 董聪朝, 钟俊, 吴凯丰, 丁勇

1830

MXene负载的非贵金属单原子催化剂催化CO氧化反应
朱纯, 梁锦霞, 王阳刚, 李隽

1842

超薄三维径向结叠层高开启电位(1.15V)光阴极用于太阳能制氢
张少波, 黄辉庭, 张之杰, 冯建勇, 刘宗光, 王军转, 徐骏, 李朝升, 余林蔚, 陈坤基, 邹志刚

1851

CdS/WO₃复合材料用于光催化选择性芳香醇氧化及析氢反应
吴玉兰, 祁明雨, 谭昌龙, 唐紫蓉, 徐艺军

1860

氧官能团MXenes用于电催化合成氨的理论筛选
高怡静, 张世杰, 孙翔, 赵伟, 卓涵, 庄桂林, 王式彬, 姚子豪, 邓声威, 钟兴, 魏中哲, 王建国

1870

错位堆叠限域和微孔缺陷固定协同生长高密度原子级Fe-N₄氧还原活性位点
陈梦慧, 陈永婷, 杨智力, 罗进, 蔡佳琳, 容忠言, 张久俊, 陈胜利, 张世明

1879

MFI/MEL分子筛催化甲醇制烯烃反应关键参数的集成方法
Chuncheng Liu, Evgeny A. Uslamin, Sophie H. van Vreeswijk, Irina Yarulina, Swapna Ganapathy, Bert M. Weckhuysen, Freek Kapteijn, Evgeny A. Pidko

1894

利用*N*-羟基邻苯二甲酰亚胺研究可见光照射下Nb₂O₅中高度扭曲的NbO₆单元作为电子转移位点
苏凯艺, 张超锋, 王业红, 张健, 郭强, 高著衍, 王峰

1906

TM@Ti₂CT_x电催化还原CO₂: 官能团诱导电子轨道重构与电荷转移
李能, 彭嘉禾, 史祖皓, 张鹏, 李鑫

1918

工况表征方法揭示燃料电池催化层中过氧化氢浓度
邱春禹, 万里洋, 王宇成, Muhammad Rauf, 洪宇浩, 袁家寅, 周志有, 孙世刚

1927

用于过氧单硫酸盐活化的超耐用氟化V₂AIC
李超, 宋晨杰, 李慧, 叶立群, 徐怡雪, 黄应平, 聂工哲, 张如梦, 刘维, 黄妞, 王保强, 马天翼

1937

在Ti₃C₂T_x MXene上原位生长2D TiO₂纳米片的异质结构用于改善电催化氮气还原
钱秀, 魏艳娇, 孙梦洁, 韩野, 张晓俐, 田健, 邵敏华


1945

介孔高硅Y沸石的合成及催化裂化性能
崔文浩, 朱大丽, 谭涓, 陈南, 樊栋, 王娟, 韩晶峰, 王林英, 田鹏, 刘中民

1955

不对称配位磷酸钴铵增强电催化水氧化反应性能
齐静, 陈明星, 张伟, 曹睿

英文全文电子版(国际版)由Elsevier出版社在ScienceDirect上出版
<https://www.sciencedirect.com/journal/chinese-journal-of-catalysis>
www.cjccatal.com
在线投审稿网址
<https://mc03.manuscriptcentral.com/cjccatal>

available at www.sciencedirect.comjournal homepage: www.sciencedirect.com/journal/chinese-journal-of-catalysis

Special column on catalytic conversion of CO₂
 Guest Editors: Yanqiang Huang, Da-Gang Yu, Guoxiong Wang

Chinese Journal of Catalysis

Graphical Contents

Special column on catalytic conversion of CO₂

Chin. J. Catal., 2022, 43: 1545–1546 doi: 10.1016/S1872-2067(22)64112-3 [Editorial]

Preface to special column on catalytic conversion of CO₂

Yanqiang Huang, Da-Gang Yu, Guoxiong Wang (Guest Editors)

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; Sichuan University




Chin. J. Catal., 2022, 43: 1547–1597 doi: 10.1016/S1872-2067(21)64000-7 [Review]

Single atom-based catalysts for electrochemical CO₂ reduction

Qian Sun, Chen Jia, Yong Zhao, Chuan Zhao *

The University of New South Wales, Australia




This review introduces the synthesis of SACs, summarizes their CO₂RR applications, discusses the characterization techniques such as *in situ/operando* skills to reveal the in-depth catalytic mechanism and provides future directions of SACs and molecular catalysts.

Chin. J. Catal., 2022, 43: 1598–1617 doi: 10.1016/S1872-2067(21)64029-9 [Review]

Recent advances in fixation of CO₂ into organic carbamates through multicomponent reaction strategies

Lu Wang, Chaorong Qi *, Wenfang Xiong, Huanfeng Jiang *
South China University of Technology

This review summarizes the recent advances in the synthesis of organic carbamates from CO₂ through multicomponent reaction strategies. Future perspectives and challenges in incorporating CO₂ into carbamates are also presented.




Chin. J. Catal., 2022, 43: 1618–1633 doi: 10.1016/S1872-2067(21)63965-7 [Review]

Tuning strategies and structure effects of electrocatalysts for carbon dioxide reduction reaction

Cong Liu, Xuanhao Mei, Ce Han, Xue Gong, Ping Song *, Weilin Xu *
Changchun Institute of Applied Chemistry, Chinese Academy of Sciences;
University of Science and Technology of China

An overview of catalysts tuning strategies has been provided, including electrochemical methods, nanocrystal synthesis, and a series of structure effects such as tandem catalysis, synergistic effects, and confinement catalysis.




Chin. J. Catal., 2022, 43: 1634–1641 doi: 10.1016/S1872-2067(21)63967-0 [Review]


Progresses on carbon dioxide electroreduction into methane

Han Zheng, Zhengwu Yang, Xiangdong Kong, Zhigang Geng *, Jie Zeng *
University of Science and Technology of China

We summarized the recent progresses on CO₂ electroreduction into CH₄ from the perspective of catalyst design strategies including construction of sub-nanometer catalytic sites, modulation of interfaces, *in-situ* structural evolution, and engineering of tandem catalysts.



Chin. J. Catal., 2022, 43: 1642–1651 doi: 10.1016/S1872-2067(21)63848-2 [Article]

Efficient hydrocarboxylation of alkynes based on carbodiimide-regulated *in situ* CO generation from HCOOH: An alternative indirect utilization of CO₂Shu-Mei Xia, Zhi-Wen Yang, Kai-Hong Chen, Ning Wang, Liang-Nian He *
Nankai University


- use carbodiimide as activator for hydrocarboxylation of alkynes
- high TON value up to 900
- no toxic CO and the indirect use of CO₂
- first synthesis of 2,2'-(1,4-phenylene)diacrylic acid
- high chem-, regio- and stereoselectively
- good results in gram scale experiment

We provide a new strategy for using carbodiimide as dehydrant to realize highly chem-, regio- and stereoselective Pd-catalyzed hydrocarboxylation of alkynes with HCOOH as both CO source and hydrogen donor, as a protocol for indirect utilization of CO₂.

Chin. J. Catal., 2022, 43: 1652–1656 doi: 10.1016/S1872-2067(21)63956-6 [Communication]

Photo-catalyzed sequential dearomatization/carboxylation of benzyl *o*-halogenated aryl ether with CO₂ leading to spirocyclic carboxylic acidsYaping Yi, Chanjuan Xi *
Tsinghua University; Nankai University


An effective method to synthesize spirocyclic carboxylic acids from benzyl *o*-halogenated aryl ether and CO₂ is developed with mild conditions, high selectivity, ideal yields and good functional group tolerance.



Chin. J. Catal., 2022, 43: 1657–1666 doi: 10.1016/S1872-2067(21)64010-X [Article]

S-Scheme 2D/2D Bi₂MoO₆/BiOI van der Waals heterojunction for CO₂ photoreductionZhongliao Wang, Bei Cheng, Liuyang Zhang *, Jiaguo Yu *, Youji Li, S. Wageh, Ahmed A. Al-Ghamdi
Wuhan University of Technology, China;
China University of Geosciences, China;
Jishou University, China;
King Abdulaziz University, Saudi Arabia


The S-scheme 2D/2D Bi₂MoO₆/BiOI van der Waals heterojunction accelerates charge separation, enhances the redox capability of photoexcited electron-hole pairs, and decreases the energy barrier of the CO₂ photoreduction reaction, contributing to a high CO and CH₄ yield rate.



Chin. J. Catal., 2022, 43: 1667–1673 doi: 10.1016/S1872-2067(21)63859-7 [Communication]

Visible-light photoredox-catalyzed carboxylation of benzyl halides with CO₂: Mild and transition-metal-free

Ke Jing, Ming-Kai Wei, Si-Shun Yan, Li-Li Liao, Ya-Nan Niu, Shu-Ping Luo, Bo Yu *, Da-Gang Yu *
Sichuan University; Zhejiang University of Technology




Visible-light photoredox-catalyzed carboxylation of benzyl halides with CO₂ was developed using organic dyes as photocatalysts under mild and transition-metal-free conditions, providing an efficient approach for assembling valuable aryl acetic acids.

Chin. J. Catal., 2022, 43: 1674–1679 doi: 10.1016/S1872-2067(21)63943-8 [Communication]

Regulating surface In–O in In@InO_x core-shell nanoparticles for boosting electrocatalytic CO₂ reduction to formate

Yan Yang, Jiaju Fu, Tang Tang, Shuai Niu, Li-Bing Zhang, Jianan Zhang, Jin-Song Hu *
Institute of Chemistry, Chinese Academy of Sciences; Zhengzhou University; University of Chinese Academy of Sciences

A facile air calcination and *in situ* reconstruction strategy is developed to regulate the surface In–O sites in core-shell In@InO_x catalysts for revealing its effects on electrocatalytic CO₂RR.




Chin. J. Catal., 2022, 43: 1680–1686 doi: 10.1016/S1872-2067(21)63999-2 [Communication]

Regulating Pd-catalysis for the electrocatalytic CO₂ reduction to formate via intermetallic PdBi nanosheets

Linfeng Xie, Xuan Liu, Fanyang Huang, Jiashun Liang, Jianyun Liu, Tanyuan Wang, Liming Yang, Ruiguo Cao, Qing Li *
Huazhong University of Science and Technology; University of Science and Technology of China

Intermetallic PdBi nanosheet with ordered hexagonal phase is developed as electrocatalyst for selective reduction of CO₂ to formate. The ordered-PdBi has a lower free energy barrier of *OCHO and higher energy barrier of *H than disordered-PdBi, enhancing the activity of formate production.




Chin. J. Catal., 2022, 43: 1687–1696 doi: 10.1016/S1872-2067(21)63970-0 [Article]

Metal organic framework-Ionic liquid hybrid catalysts for the selective electrochemical reduction of CO₂ to CH₄

Ernest Pahuyo Delmo, Yian Wang, Jing Wang, Shangqian Zhu, Tiehuai Li, Xueping Qin, Yibo Tian, Qinglan Zhao, Juhee Jang, Yinu Wang, Meng Gu, Lili Zhang *, Minhua Shao *
The Hong Kong University of Science and Technology; Southern University of Science and Technology; Huaiyin Normal University


MOF-RTIL hybrid electrocatalysts are shown to selectively reduce CO₂ to CH₄ with a maximum FE of 65%. This study illustrates a novel approach of embedding co-catalysts within porous frameworks to improve CO₂RR electrocatalysis.



Chin. J. Catal., 2022, 43: 1697–1702 doi: 10.1016/S1872-2067(21)64006-8 [Article]

Synergetic effect of nitrogen-doped carbon catalysts for high-efficiency electrochemical CO₂ reduction

Chuhao Liu, Yue Wu, Jinjie Fang, Ke Yu, Hui Li, Wenjun He, Weng-Chon Cheong, Shoujie Liu, Zheng Chen, Jing Dong*, Chen Chen*
Tsinghua University; Nanjing Tech University; Beijing University of Chemical Technology; University of Macau; Anhui Normal University




Metal-free nitrogen-doped carbon catalysts (mf-NCs) with well-defined active N-species sites were synthesized as robust catalysts for CO₂RR. Graphitic N and pyridinic N can synergistically improve CO₂-to-CO selectivity.

Chin. J. Catal., 2022, 43: 1703–1709 doi: 10.1016/S1872-2067(21)63988-8 [Article]

Electroreduction of air-level CO₂ with high conversion efficiency

Yangshen Chen, Miao Kan, Shuai Yan, Junbo Zhang, Kunhao Liu, Yaqin Yan, Anxiang Guan, Ximeng Lv, Linping Qian, Gengfeng Zheng*
Fudan University




This paper realizes the direct electroreduction of 0.03% or 0.04% CO₂ in a homemade gas-phase electrochemical reactor using a low-cost copper (Cu) or nanoscale copper (nano-Cu) catalyst with high CO₂ conversion efficiency.

Chin. J. Catal., 2022, 43: 1710–1718 doi: 10.1016/S1872-2067(21)63960-8 [Article]

Highly dispersed nickel species on iron-based perovskite for CO₂ electrolysis in solid oxide electrolysis cell

Yingjie Zhou, Tianfu Liu, Yuefeng Song, Houfu Lv, Qingxue Liu, Na Ta, Xiaomin Zhang*, Guoxiong Wang*
Donghua University; Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences



Highly dispersed nickel species with low loading on perovskite can alter its electronic structure and facilitate the oxygen vacancy formation, thus greatly promote the CO₂ electrolysis.


Reviews

Chin. J. Catal., 2022, 43: 1719–1748 doi: 10.1016/S1872-2067(21)63994-3

Environmentally-friendly carbon nanomaterials for photocatalytic hydrogen production

Sheng Xiong, Rongdi Tang, Daoxin Gong, Yaocheng Deng *, Jiangfu Zheng, Ling Li, Zhanpeng Zhou, Lihua Yang, Long Su
Hunan Agricultural University


This review introduced the CNMs, including carbon dots, fullerenes, carbon nanotubes, carbon nanofibers, graphene, and graphdiyne. CNMs can provide abundant adsorption and active sites, charge separation, transport channels, and can be used as photocatalysts, co-catalysts and photosensitizers.



Chin. J. Catal., 2022, 43: 1749–1760 doi: 10.1016/S1872-2067(21)64002-0

Toward modular construction of cell-free multienzyme systems

Yinchen Zhang, Ning Nie, Yifei Zhang *
Beijing University of Chemical Technology




Modulization has shown advantages in the construction and optimization of cell-free multienzyme systems. Here we introduce typical natural and synthetic multienzyme modules at and across multiple levels.

Chin. J. Catal., 2022, 43: 1761–1773 doi: 10.1016/S1872-2067(21)64001-9


Photoelectrochemical nitrogen reduction: A step toward achieving sustainable ammonia synthesis

Liqun Wang, Xiao Yan, Wenping Si, Daolan Liu, Xinggong Hou, Dejun Li *, Feng Hou *, Shi Xue Dou, Ji Liang *
Tianjin Normal University, China;
Shenzhen Institute of Information Technology, China;
Tianjin University, China;
University of Wollongong, Australia

The photoelectrochemical NRR is an energy-efficient and environmentally friendly route for sustainable ammonia synthesis. In this review, the fundamental principles and the latest progress in research into photocathode materials of PEC-NRR are described.



Chin. J. Catal., 2022, 43: 1774–1804 doi: 10.1016/S1872-2067(22)64105-6

Photodeposition of earth-abundant cocatalysts in photocatalytic water splitting: Methods, functions, and mechanismsHui Zhao, Qinyi Mao, Liang Jian, Yuming Dong*, Yongfa Zhu*
Jiangnan University; Tsinghua University


This review provides a detailed insight into the photodeposition of earth-abundant cocatalysts onto semiconductors for photocatalytic water splitting, including the deposition methods, catalytic performance, and functional mechanisms.

Communications

Chin. J. Catal., 2022, 43: 1805–1811 doi: 10.1016/S1872-2067(22)64101-9

Aromatic bromination with hydrogen production on organic-inorganic hybrid perovskite-based photocatalysts under visible light irradiationYanfei Zhang, Hong Wang, Yan Liu, Can Li*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences;
University of Chinese Academy of Sciences


Organic-inorganic hybrid perovskite MAPbBr₃, hybridizing with Pt-Ta₂O₅ and PEDOT:PSS, can achieve aromatic bromination along hydrogen production with high yield and selectivity using HBr as the bromine source under visible light irradiation.



Chin. J. Catal., 2022, 43: 1812–1817 doi: 10.1016/S1872-2067(21)64051-2

The synthesis of tetracyclic coumarins via decarboxylative asymmetric [4+2] cycloadditions enabled by Pd(0)/Cu(I) synergistic catalysisKai Wang, Xiangfeng Lin, Qian Li, Yan Liu*, Can Li*
Dalian Institute of Chemical Physics, Chinese Academy of Sciences;
University of Chinese Academy of Sciences

A highly efficient asymmetric decarboxylative [4+2] cycloaddition of 3-cyanocoumarins with vinyl benzoxazinones was developed, affording the coumarin-derived condensed rings bearing three continuous stereocenters in high yields with excellent diastereoselectivities (>20/1 d.r.) and enantioselectivities (up to 99% ee).



Articles

Chin. J. Catal., 2022, 43: 1818–1829 doi: 10.1016/S1872-2067(21)64009-3

Energy funneling and charge separation in CdS modified with dual cocatalysts for enhanced H₂ generation

Meiyu Zhang, Chaochao Qin, Wanjun Sun, Congzhao Dong, Jun Zhong, Kaifeng Wu, Yong Ding *

Lanzhou University;


Dalian Institute of Chemical Physics, Chinese Academy of Sciences;

Henan Normal University;

Soochow University;

Lanzhou Institute of Chemical Physics, Chinese Academy of Sciences

For CdS/MBI/MCoA, the MBI and MCoA molecules selectively extracted photogenerated holes and electrons from CdS nanorods rapidly. Additionally, the MBI molecules facilitated energy funneling from thin to thick rods, endowing CdS/MBI/MCoA elevated photocatalytic H₂ evolution.



Chin. J. Catal., 2022, 43: 1830–1841 doi: 10.1016/S1872-2067(21)64027-5

Non-noble metal single-atom catalyst with MXene support: Fe₁/Ti₂CO₂ for CO oxidation


Chun Zhu, Jin-Xia Liang*, Yang-Gang Wang, Jun Li *

Guizhou University;

Southern University of Science and Technology;

Tsinghua University

MXene supported non-noble metal single-atom catalyst Fe₁/Ti₂CO₂ is found to exhibit excellent activity for low-temperature CO oxidation through first-principles computational modelling.




Chin. J. Catal., 2022, 43: 1842–1850 doi: 10.1016/S1872-2067(21)64046-9

Ultrathin 3D radial tandem-junction photocathode with a high onset potential of 1.15 V for solar hydrogen production

Shaobo Zhang, Huiting Huang, Zhijie Zhang, Jianyong Feng, Zongguang Liu, Junzhuan Wang, Jun Xu, Zhaosheng Li *, Linwei Yu *, Kunji Chen, Zhigang Zou

Nanjing University




A radial tandem junction thin-film photocathode with a V_{onset} of 1.15 V vs. RHE has been demonstrated directly on VLS-grown SiNWs for the first time, using a-Si:H (~50 nm) as the outer absorber layer and a-SiGe:H (~40 nm) as the inner layer.

Chin. J. Catal., 2022, 43: 1851–1859 doi: 10.1016/S1872-2067(21)63989-X

Photocatalytic selective oxidation of aromatic alcohols coupled with hydrogen evolution over CdS/WO₃ composites

Yu-Lan Wu, Ming-Yu Qi, Chang-Long Tan, Zi-Rong Tang*, Yi-Jun Xu*
Fuzhou University




The photoexcited charge carriers can be effectively separated by the construction of a direct Z-scheme system. Under light irradiation, photocatalytic selective oxidation of aromatic alcohols to aromatic aldehydes coupled with hydrogen evolution are simultaneously realized on the Z-scheme CdS/WO₃ photocatalyst, making full use of the synergistic potential of the combined photoredox system.

Chin. J. Catal., 2022, 43: 1860–1869 doi: 10.1016/S1872-2067(21)64011-1

Computational screening of O-functional MXenes for electrocatalytic ammonia synthesis

Yijing Gao, Shijie Zhang, Xiang Sun, Wei Zhao, Han Zhuo, Guilin Zhuang, Shibin Wang, Zihao Yao, Shengwei Deng, Xing Zhong, Zhongzhe Wei, Jian-guo Wang*
Zhejiang Normal University; Zhejiang University of Technology




A three-step screening method (including stability, activity, and selectivity screening) was applied to develop NRR catalysts from O-functional MXenes, and LASSO regression was applied to describe the original NRR performance.

Chin. J. Catal., 2022, 43: 1870–1878 doi: 10.1016/S1872-2067(21)63992-X

Synergy of staggered stacking confinement and microporous defect fixation for high-density atomic Fe^{II}-N₄ oxygen reduction active sites

Menghui Chen, Yongting Chen, Zhili Yang, Jin Luo, Jialin Cai, Joey Chung-Yen Jung, Jiujun Zhang, Shengli Chen*, Shiming Zhang*
Shanghai University; Wuhan University



A synergistic modulation strategy of phthalocyanine (Pc) for spatial-locations confinement and Zn-salts for active-sites fixation has been proposed to overwhelmingly grow atomic Fe^{II}-N₄ for the high ORR performance Fe-N_x-C catalyst.


Chin. J. Catal., 2022, 43: 1879–1893 doi: 10.1016/S1872-2067(21)63990-6

An integrated approach to the key parameters in methanol-to-olefins reaction catalyzed by MFI/MEL zeolite materials

Chuncheng Liu, Evgeny A. Uslamin, Sophie H. van Vreeswijk, Irina Yarulina, Swapna Ganapathy, Bert M. Weckhuysen, Freek Kapteijn*, Evgeny A. Pidko*

Delft University of Technology, the Netherlands;

Process Research and Chemical Engineering, Germany; Utrecht University, the Netherlands




Al enrichment in the channels is of primary importance to determine propylene production in MTO. Catalyst deactivation is rather complex, which is controlled by Al distribution, crystal size, topology and Al_{pair}.

Chin. J. Catal., 2022, 43: 1894–1905 doi: 10.1016/S1872-2067(21)64026-3

Unveiling the highly disordered NbO₆ units as electron-transfer sites in Nb₂O₅ photocatalysis with *N*-hydroxyphthalimide under visible light irradiation

Kaiyi Su, Chaofeng Zhang*, Yehong Wang, Jian Zhang, Qiang Guo, Zhuyan Gao, Feng Wang*

Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences




Using *N*-hydroxyphthalimide as probe molecule, highly disordered NbO₆ units act as the electron-transfer sites on Nb₂O₅ under visible light irradiation, which can be conducive to a deep understanding of NbO_x units in photocatalysis.

Chin. J. Catal., 2022, 43: 1906–1917 doi: 10.1016/S1872-2067(21)64018-4

Charge transfer and orbital reconstruction of non-noble transition metal single-atoms anchored on Ti₂CT_x-MXenes for highly selective CO₂ electrochemical reduction

Neng Li *, Jiahe Peng, Zuhao Shi, Peng Zhang, Xin Li *

Wuhan University of Technology; Zhengzhou University; South China Agricultural University




Oxygen surface group can provide adequate electrons in CO₂ activation. The charge transfer and orbital reconstruction between CO₂ and TM@Ti₂CO₂ leads to the activation of CO₂, which does not happen at -S surface group.

Chin. J. Catal., 2022, 43: 1918–1926 doi: 10.1016/S1872-2067(21)63993-1

Revealing the concentration of hydrogen peroxide in fuel cell catalyst layers by an *in-operando* approachChun-Yu Qiu, Li-yang Wan, Yu-Cheng Wang *, Muhammad Rauf, Yu-Hao Hong, Jia-yin Yuan, Zhi-You Zhou *, Shi-Gang Sun
Xiamen University, China; Shenzhen University, China; Stockholm University, Sweden


The *in-operando* concentration of H₂O₂ in Fe/N/C cathode catalyst layer is one order of magnitude higher than that under aqueous electrodes test conditions.



Chin. J. Catal., 2022, 43: 1927–1936 doi: 10.1016/S1872-2067(21)64050-0

Ultradurable fluorinated V₂AlC for peroxydisulfate activation in organic pollutant degradation processesChao Li, Chenjie Song, Hui Li, Liqun Ye *, Yixue Xu, Yingping Huang, Gongzhe Nie, Rumeng Zhang, Wei Liu, Niu Huang, Po Keung Wong, Tianyi Ma *
China Three Gorges University, China; Hubei Three Gorges Laboratory, China; Swinburne University of Technology, Australia; The Chinese University of Hong Kong, China

F-V₂AlC showed higher activity and better reusability than V₂O₃ for PMS activation with extremely low ion leakage. And mechanism investigation showed that the main reactive oxygen specie was ¹O₂, which induced by the two-dimensional confinement effect. In addition, the introduction of F changed the adsorption mode of TC over catalyst, thereby changing the degradation path.




Chin. J. Catal., 2022, 43: 1937–1944 doi: 10.1016/S1872-2067(21)64020-2

Heterostructuring 2D TiO₂ nanosheets *in situ* grown on Ti₃C₂T_x MXene to improve the electrocatalytic nitrogen reduction

Xiu Qian, Yanjiao Wei, Mengjie Sun, Ye Han *, Xiaoli Zhang, Jian Tian *, Minhua Shao *

Shandong University of Science and Technology;
The Hong Kong University of Science and Technology;
Zhengzhou University


TiO₂/Ti₃C₂T_x MXene was prepared through the *in situ* growth of TiO₂ nanosheets (NSs) on extremely conductive Ti₃C₂T_x MXene, which exhibited an excellent electrochemical activity for NRR (NH₃ yield of 44.17 μg h⁻¹ mg⁻¹_{cat.}).



Chin. J. Catal., 2022, 43: 1945–1954 doi: 10.1016/S1872-2067(21)64043-3

Synthesis of mesoporous high-silica zeolite Y and their catalytic cracking performance

Wenhao Cui, Dali Zhu, Juan Tan, Nan Chen, Dong Fan, Juan Wang, Jingfeng Han, Linying Wang *, Peng Tian *, Zhongmin Liu
Dalian Institute of Chemical Physics, Chinese Academy of Sciences; University of Chinese Academy of Sciences; Dalian University of Technology



Mesoporous high-silica zeolite Y with high (hydro)thermal stability has been successfully synthesized, which possesses larger acid concentration and stronger acid strength than USY, and exhibits superior catalytic performance in the cracking of hydrocarbons and industrial heavy oil.

Chin. J. Catal., 2022, 43: 1955–1962 doi: 10.1016/S1872-2067(21)64035-4

Ammonium cobalt phosphate with asymmetric coordination sites for enhanced electrocatalytic water oxidation

Jing Qi, Mingxing Chen, Wei Zhang *, Rui Cao *
Henan Normal University; Shaanxi Normal University

Two cobalt phosphates with distinct Co symmetry were obtained to explore the effect of coordination structure on the electrocatalytic oxygen evolution reaction. Asymmetric Co geometry can accelerate surface reconstruction to enhance OER activity.

