


ISSN 0567-7351
CN 31-1320/O6
CODEN HHPA4
<http://sioc-journal.cn>



化学学报

ACTA CHIMICA SINICA

2017 第75卷 第8期 Vol. 75 No. 8



ISSN 0567-7351



中国化学会
中国科学院上海有机化学研究所

主办

万方数据

化学学报

Acta Chimica Sinica

(Huaxue Xuebao)

第 75 卷 第 8 期 2017 年 8 月 15 日

目 次

综述

- 从氮气直接合成含氮有机化合物 李嘉鹏, 殷剑昊, 俞超, 张文雄, 席振峰*, 化学学报, **2017**, 75(8), 733-743
- 直接三氟甲磺基化试剂及方法的研究进展 张盼盼, 吕龙, 沈其龙*, 化学学报, **2017**, 75(8), 744-769
- 光控释药型药物递送系统的研究进展 张留伟, 钱明, 王静云*, 化学学报, **2017**, 75(8), 770-782

研究通讯

- 铜(II)催化的吡啶与四取代 D-A 环丙烷[3+2]开环/环化反应 严文广, 王盼, 王丽佳, 孙秀丽*, 唐勇*, 化学学报, **2017**, 75(8), 783-787
- 酸性体系 V 催化木质素 β -O-4 模型物 C—C 键高选择性切断 刘新鑫, 严龙, 傅尧*, 化学学报, **2017**, 75(8), 788-793
- 铜催化苯乙烯不对称硼胺化反应 张涌灵, 王敏, 曹鹏, 廖建*, 化学学报, **2017**, 75(8), 794-797

研究论文


- Cortistatin 类型天然产物的不对称形式全合成: 金催化串联 Semi-Pinacol 重排反应策略 顾月青, 袁浩, 傅俊凯*, 龚建贤*, 杨震*, 化学学报, **2017**, 75(8), 798-807
- 基于萘并二酰亚胺的胺基功能化聚合物的三组分一锅法合成及其在聚合物太阳能电池中的应用 贾涛, 郑楠楠, 蔡万清, 应磊, 黄飞*, 化学学报, **2017**, 75(8), 808-818
- 芳香胺盐酸盐/硼烷体系催化的氢胺化/还原反应研究 张露文, 温志国, Borzov, Maxim, 聂万丽*, 化学学报, **2017**, 75(8), 819-823
- 有机胺盐/硼烷体系与炔烃的硼氢化加成反应机理研究 孙国峰, 苏敏, 方洁, Borzov, Maxim, 聂万丽*, 化学学报, **2017**, 75(8), 824-830

* 通信联系人.

On the cover: This review mainly summarizes the progress in the field of direct transformation of molecular nitrogen to nitrogen-containing organic compounds by using transition metal complexes. The activation and transformation of dinitrogen (nitrogen fixation) under mild conditions has been a great challenge, which requires combined efforts of chemists from all over the world. [Xi, Zhenfeng *et al.* on page 733-743.]



On the back cover: A bisoxazoline/Cu(SbF₆)₂ catalyzed stereoselective [3+2] annulation of indoles with 1,1,2,2-tetrasubstituted donor-acceptor cyclopropanes was presented. This annulation method furnished the C2, C3-fused indolines bearing three continuous quaternary stereocentres on the newly built cyclopentane ring, which are widely existed as a core motif in a plenty of natural products and biologically active compounds. [Tang, Yong *et al.* on page 783-787.]




Review

Direct Transformation of N₂ to N-Containing Organic Compounds

Li, Jiapeng; Yin, Jianhao; Yu, Chao; Zhang, Wenxiong; Xi, Zhenfeng*


Acta Chim. Sinica **2017**, 75(8), 733-743



Recent Progress on Direct Trifluoromethylthiolating Reagents and Methods


Zhang, Panpan; Lu, Long; Shen, Qilong*

Acta Chim. Sinica **2017**, 75(8), 744-769



Recent progress on the development of trifluoromethylthiolating reagents and methods for the direct introduction of the trifluoromethylthio group was reviewed.

Progress in Research of Photo-controlled Drug Delivery Systems




Zhang, Liuwei; Qian, Ming; Wang, Jingyun*
Acta Chim. Sinica **2017**, 75(8), 770-782

Many photo-controlled drug delivery systems based on different photo-responsive groups for the precise release of drugs are summarized in this review. Photo-controlled drug delivery systems provide a new strategy for the precise treatment of tumors.

Communication

Copper Catalyzed [3+2] Annulation of Indoles with 1,1,2,2-Tetrasubstituted Donor-Acceptor Cyclopropanes




Yan, Weng-Guang; Wang, Pan; Wang, Lijia;
 Sun, Xiu-Li*; Tang, Yong*

Acta Chim. Sinica **2017**, 75(8), 783-787

A first Lewis acid catalyzed [3+2] annulation of indoles with 1,1,2,2-tetrasubstituted D-A cyclopropanes was reported, providing an easy access to a variety of C2, C3-fused indolines, bearing three quaternary stereocentres on the newly built cyclopentane ring.

Lignin C—C Bond's Cleavage by Vanadium Catalyzed with High Selectivity in Acid Environment




Liu, Xinxin; Yan, Long; Fu, Yao*

Acta Chim. Sinica **2017**, 75(8), 788-793

The process of pre-oxidized lignin β -O-4 model compound converted to benzoic acid and phenol has been studied. The β -O-4 structure would be degraded through C—C bonds' cleavage efficiently over the catalyst of NH_4VO_3 . It has been proved that oxovanadium(V) cations in the system were the core for the catalyzed process and recycle via oxovanadium(IV) cations.

Copper-Catalyzed Enantioselective Aminoboration of Styrenes with Chiral Sulfoxide Phosphine Ligand




Zhang, Yongling; Wang, Min; Cao, Peng; Liao, Jian*

Acta Chim. Sinica 2017, 75(8), 794-797

A Cu-catalyzed enantioselective aminoboration of styrenes by using chiral sulfoxide-phosphine (SOP) ligands was developed, chiral β -aminoalkylboranes were obtained with satisfied yields and *ee* values, which can be readily transferred to valuable β -hydroxylalkylamines.

Article

Asymmetric Formal Synthesis of Cortistatins via a Gold-Catalyzed Semi-Pinacol Rearrangement Strategy




Gu, Yueqing; Yuan, Hao; Fu, Junkai*; Gong, Jianxian*; Yang, Zhen*

Acta Chim. Sinica 2017, 75(8), 798-807

Full details about our efforts towards the formal synthesis of cortistatins were described herein. This route is featured with a novel gold-catalyzed cascade reaction involving intramolecular nucleophilic addition of hydroxyl group to the carbon-carbon triple bond, followed by an oxonium ions initiated semi-pinacol-type 1,2-migration to construct the key oxabicyclo[3.2.1]octane skeleton.

Naphthalene Diimide-Based Polymers Consisting of Amino Alkyl Side Groups: Three-Component One-Pot Polymerization and Their Application in Polymer Solar Cells




Jia, Tao; Zheng, Nannan; Cai, Wanqing; Ying, Lei; Huang, Fei*

Acta Chim. Sinica 2017, 75(8), 808-818

A series of naphthalene diimide-based amino-functionalized polymers **P1**~**P4** were developed via three-component polymerization (TCP) of diynes, dialdehydes and dibenzylamine. The influence of chemical environment of amine groups on the electrode modification capability and self-doping behavior of the resulting polymers was explored. The **P1** can be utilized as the cathode interlayer for polymer solar cells and the resulting device exhibited a significantly improved performance. Our results indicate that TCP is an effective strategy for the development of multi-functional polymer materials.

Research of B(C₆F₅)₃/Aromatic Ammonium Chloride Systems Catalyzed Hydroamination/Reduction Reaction



Zhang, Luwen; Wen, Zhiguo; Borzov, Maxim; Nie, Wanli*


Acta Chim. Sinica 2017, 75(8), 819-823

A new method of an one-pot hydroamination/reduction reaction of terminal alkyne with aromatic amines catalyzed by the B(C₆F₅)₃/aromatic ammonium chloride systems was described. Hydrosilane has been chosen as reducing reagent instead of hydrogen.

Research of the Stereoselectivity and Mechanism of the Hydroboration Reaction Between $B(C_6F_5)_3$ /Ammonium Chloride Systems with Terminal Alkyne

Sun, Guofeng; Su, Min; Fang, Jie; Borzov, Maxim; Nie, Wanli*

Acta Chim. Sinica **2017**, 75(8), 824-830



$R''_2NH = t\text{-BuNH}_2, \text{CyNH}_2, i\text{-Pr}_2\text{NH}, \text{Et}_2\text{NH}, \text{TMP}, \text{Et}_3\text{NH}, n\text{-Bu}_4\text{N}, \text{etc.}$

This paper describes a series of reactions between $B(C_6F_5)_3$ /ammonium chloride systems with terminal alkynes and the characteristics of the Z-stereoselective hydroboration products. The mechanism of the stereoselectivity has been discussed and found that the ammonium hydroborate $[R_2NH_2][HB(C_6F_5)_3]$ intermediates could not react with the alkynes alone. Trace amount of the Lewis acid $B(C_6F_5)_3$ is necessary to firstly activate the alkynes. And the weak interaction between the Cl ion and $B(C_6F_5)_3$ plays also an important role on these stereoselective hydroboration reactions.




ISSN 0567-7351
CN 31-1320/O6
CODEN HHHPA4
<http://sioc-journal.cn>


化学学报

ACTA CHIMICA SINICA



2017 第75卷 第8期 Vol. 75 No. 8






Kopsane



Vindoline



Dasyrachine



Borreverine

ISSN 0567-7351



9 770567 735172

万方数据

08>



中国化学会
中国科学院上海有机化学研究所

主办