

石油实验地质

第38卷第3期 2016年5月

目次

盆地·油藏

裂谷与被动陆缘叠合盆地的盐构造与油气成藏——以西非下刚果—刚果扇盆地和宽扎盆地为例 孙自明,何治亮(287)

陆相碎屑岩中钙质胶结物沉淀机制及油气储集意义——来自济阳拗陷孔店组和川西须家河组的对比研究 谭先锋,蒋威,吴康军,王浩,徐田堃,陈苏军,冉天(293)

川西拗陷陆相脉冲式波动“二元”层序特征研究 张庄,叶素娟,杨克明,朱宏权,杨映涛,李旻,张世华(303)

川东南寒武系娄山关群白云岩成岩作用特征 江文剑,侯明才,邢凤存,徐胜林,林良彪(311)

川东南五峰—龙马溪组页岩储层六性特征及主控因素分析 张汉荣,王强,倪楷,李春燕(320)

塔里木盆地西南拗陷东部构造特征及对油气成藏的控制 何娟,王毅,刘士林,何登发(326)

塔里木盆地麦盖提斜坡构造样式分析 史政,徐征遥(333)

美国俄克拉荷马州伍德福德页岩甜点控制因素研究 高占京,郑和荣,黄韬(340)

荆门地区页岩储层吸附特征及吸附势理论应用 岳长涛,李术元,李林玥,温海龙(346)

俄罗斯季曼—伯朝拉盆地碳酸盐岩油气藏特征分析——以 Trebs 和 Titov 油气田为例 杨士伟,史丹妮(354)

伊拉克米桑油田群 Mishrif 组 MB₂ 段碳酸盐岩储层特征及成因 张义楷,康安,闵小刚,李志明,李伟超,高慧君(360)

油气地球化学

四川盆地西部中三叠统雷口坡组烃源岩生烃潜力分析 杨克明(366)

风化作用对高演化黑色岩系有机质影响因素分析——以塔里木盆地柯坪地区玉尔吐斯组为例 陶国亮,申宝剑,腾格尔,仰云峰,徐二社,潘安阳(375)

伦坡拉盆地始新统牛堡组烃源岩成熟史 潘磊,曹强,刘一茗,李一腾,汪岩,李智泉(382)

方法·技术

核磁共振与高压压汞实验联合表征致密油储层微观孔喉分布特征 公言杰,柳少波,赵孟军,谢红兵,刘可禹(389)

深埋砂岩储层长石溶孔率定量计算的新方法及应用——以鄂尔多斯盆地陇东地区长₈储层为例 杨友运,刘喜强,孙睿(395)

油页岩不同温度原位热解物性变化核磁共振分析 李广友,马中良,郑家锡,鲍芳,郑伦举(402)

基于 X-CT 技术对黔西滇东区域煤储层物性特征的研究 熊波,刘坤,郭凯,赵广民(407)

塔里木盆地志留系碎屑岩孔隙成因类型定量分析——以顺9井柯坪塔格组为例 张永东,王恕一(413)

微米 CT 在烃源岩微观结构表征方面的应用 黄振凯,陈建平,王义军,王华建,邓春萍(418)

综合信息

中国石化无锡石油地质研究所实验地质技术之油气微生物检测技术 (封二)

《石油实验地质》征稿简则 (封三)

CONTENTS

- Salt tectonics and its relationship to hydrocarbon accumulation in salt basins with a lower rifted section and an upper continental marginal section: A case study of the Lower Congo–Congo Fan basins and the Kwanza Basin in West Africa *Sun Ziming, He Zhiliang* (287)
- Sedimentation mechanism and petroleum significance of calcareous cements in continental clastic rocks: Comparison between the Kongdian Formation in the Jiyang Depression and the Xujiahe Formation in the western Sichuan Basin *Tan Xianfeng, Jiang Wei, Wu Kangjun, Wang Hao, Xu Tiankun, Chen Sujun, Ran Tian* (293)
- LST and T-HST featured by impulse in continental sequences in the Western Sichuan Depression *Zhang Zhuang, Ye Sujuan, Yang Keming, Zhu Hongquan, Yang Yingtao, Li Min, Zhang Shihua* (303)
- Diagenetic characteristics of dolomites in the Cambrian Loushanguan Group in southeastern Sichuan Basin *Jiang Wenjian, Hou Mingcai, Xing Fengcun, Xu Shenglin, Ling Liangbiao* (311)
- Six characteristics and main controlling factors of shale reservoirs in the Wufeng–Longmaxi formations, south-eastern Sichuan Basin *Zhang Hanrong, Wang Qiang, Ni Kai, Li Chunyan* (320)
- Characteristics of fault structure and its control on hydrocarbon accumulation in the eastern part of southwestern Tarim Basin *He Juan, Wang Yi, Liu Shilin, He Dengfa* (326)
- Structural pattern of the Maigaiti Slope in the Tarim Basin *Shi Zheng, Xun Zhengyao* (333)
- Attributes of sweet spots in the Devonian Woodford shales in Oklahoma, USA *Gao Zhanjing, Zheng Herong, Huang tao* (340)
- Adsorption characteristics of shale reservoirs in the Jingmen area and application of adsorption potential theory *Yue Changtao, Li Shuyuan, Li Linyue, Wen Hailong* (346)
- Characteristics of carbonate reservoirs in the Timan-Pechora Basin in Russia: The Trebs and Titov oil fields as examples *Yang Shiwei, Shi Danni* (354)
- Characteristics and genesis of carbonate reservoirs in the Mishrif MB₂ member in the Missan oil fields, Iraq *Zhang Yikai, Kang An, Min Xiaogang, Li Zhiming, Li Weichao, Gao Huijun* (360)
- Hydrocarbon potential of source rocks in the Middle Triassic Leikoupo Formation in the Western Sichuan Depression *Yang Keming* (366)
- Weathering effects on high-maturity organic matter in a black rock series: A case study of the Yuertusi Formation in Kalpin area, Tarim Basin *Tao Guoliang, Shen Baojian, Tenger Boltsjin, Yang Yunfeng, Xu Ershe, Pan Anyang* (375)
- Maturity history of source rocks in the Eocene Niubao Formation, Lunpola Basin *Pan Lei, Cao Qiang, Liu Yiming, Li Yiteng, Wang Yan, Li Zhiquan* (382)
- Characterization of micro pore throat radius distribution in tight oil reservoirs by NMR and high pressure mercury injection *Gong Yanjie, Liu Shaobo, Zhao Mengjun, Xie Hongbing, Liu Keyu* (389)
- A new method for the calculation of secondary porosity originating from the dissolution of feldspars in deeply buried formations and its application: A case study of the Chang 8₁ Formation in Longdong area, Ordos Basin *Yang Youyun, Liu Xiqiang, Sun Rui* (395)
- NMR analysis of the physical change of oil shales during in situ pyrolysis at different temperatures *Li Guangyou, Ma Zhongliang, Zheng Jiayi, Bao Fang, Zheng Lunju* (402)
- Characterization of the physical properties of coal reservoirs in the western Guizhou and eastern Yunnan by X-ray computed tomography *Xiong Bo, Liu Kun, Guo Kai, Zhao Guangmin* (407)
- Quantitative analysis of pore genetic types in the Silurian clastic rocks in the Tarim Basin: A case study of the Kepingtage Formation in well Shun9 *Zhang Yongdong, Wang Shuyi* (413)
- Application of micron CT in the characterization of microstructure in source rocks *Huang Zhenkai, Chen Jianping, Wang Yijun, Wang Huajian, Deng Chunping* (418)