

植物生态学报

Chinese Journal of Plant Ecology

第36卷 第9期 2012年9月

Vol. 36 No. 9 September 2012



主办单位：中国科学院植物研究所
中国植物学会

Sponsors: Institute of Botany, Chinese Academy of Sciences
Botanical Society of China

植 物 生 态 学 报

Zhiwu Shengtai Xuebao

2012年9月 第36卷 第9期

目 次

研究论文	养空间异质性改变的响应
	杨秀云 韩有志 武小钢
923 长白山不同演替阶段针阔混交林群落物种 多度分布格局	973 丛枝菌根真菌通过上调根系及自身水孔蛋 白基因表达提高玉米抗旱性
闫 琰 张春雨 赵秀海	李 涛 陈保冬
935 尖峰岭热带山地雨林海南特有木本植物群 落结构	982 涝渍对3个树种生长、组织孔隙度和渗漏氧 的影响
盛大勇 庄雪影 许 涵 李意德 胡跃华	汪贵斌 曹福亮 王 媛
时雷雷 骆士寿 陈德祥 林明猷	992 基于层次-向量法分析的普通鹿蹄草品质与 生态因子的相关性
948 种子异型植物异子蓬的生殖分配和结实格局	吕振江 王冬梅 李登武
王 雷 董 鸣 黄振英	
956 芦芽山林线白杆生长季径向生长动态	综述
董满宇 江 源 杨浩春 王明昌 张文涛	1004 树干液流径向分布格局研究进展
郭媛媛	徐 飞 杨风亭 王辉民 戴晓琴
965 华北落叶松林细根生物量对土壤水分、氮营	

封面说明：长白山椴树红松林秋季景观。椴树红松林是长白山地区的典型地带性植被类型，处于群落演替的顶级阶段，主要树种有红松、紫椴、大青杨和臭松等。闫琰等研究了长白山针阔混交林群落演替进程中物种多度分布格局形成的生态学过程(本期923-934页) (赵秀海摄)。

CONTENTS

Research Articles

- 923 Species-abundance distribution patterns at different successional stages of conifer and broad-leaved mixed forest communities in Changbai Mountains, China
YAN Yan, ZHANG Chun-Yu, and ZHAO Xiu-Hai
- 935 Community structure of endemic woody plants in tropical montane rainforest of Jianfengling, Hainan Island, China
SHENG Da-Yong, ZHUANG Xue-Ying, XU Han, LI Yi-De, HU Yue-Hua, SHI Lei-Lei, LUO Tu-Shou, CHEN De-Xiang, and LIN Ming-Xian
- 948 Reproductive allocation and fruit-set pattern of seed-heteromorphic plant *Suaeda aralocaspica*
WANG Lei, DONG Ming, and HUANG Zhen-Ying
- 956 Dynamics of stem radial growth of *Picea meyeri* during the growing season at the treeline of Luya Mountain, China
DONG Man-Yu, JIANG Yuan, YANG Hao-Chun, WANG Ming-Chang, ZHANG Wen-Tao, and GUO Yuan-Yuan
- 965 Response of fine root biomass to changes in spatial heterogeneity of soil moisture and nitrogen in *Larix principis-rupprechtii* forest
YANG Xiu-Yun, HAN You-Zhi, and WU Xiao-Gang
- 973 Arbuscular mycorrhizal fungi improving drought tolerance of maize plants by up-regulation of aquaporin gene expressions in roots and the fungi themselves
LI Tao and CHEN Bao-Dong
- 982 Effects of waterlogging on growth, porosity and radial oxygen loss of three tree species
WANG Gui-Bin, CAO Fu-Liang, and WANG Yuan
- 992 Correlation between quality of *Pyrola decorata* and its ecological factors based on hierarchy-vector analysis
LÜ Zhen-Jiang, WANG Dong-Mei, and LI Deng-Wu

Review

- 1004 Review of advances in radial patterns of stem sap flow
XU Fei, YANG Feng-Ting, WANG Hui-Min, and DAI Xiao-Qin

Cover illustration: The autumn landscape of *Tilia amurensis*-*Pinus koraiensis* mixed forest in Changbai Mountain. *Tilia amurensis*-*Pinus koraiensis* mixed forest is the typical zonal vegetation type in the Changbai Mountain, which is in the climax of the community succession. The major tree species include *Pinus koraiensis*, *Tilia amurensis*, *Populus ussuriensis* and *Abies nephrolepis*. Yan *et al.* investigated the ecological processes that shape species-abundance distribution pattern in the community succession process of the conifer and broad-leaved mixed forest in Changbai Mountain (Pages 923–934 of this issue) (Photographed by ZHAO Xiu-Hai).